

Asymptotics of a vanishing period : the quotient themes of a given fresco.

Daniel Barlet*

13/01/11

Abstract

In this paper we introduce the word "fresco" to denote a $[\lambda]$ -primitive monogenic geometric (a,b)-module. The study of this "basic object" (generalized Brieskorn module with one generator) which corresponds to the minimal filtered (regular) differential equation satisfied by a relative de Rham cohomology class, began in [B.09] where the first structure theorems are proved. Then in [B.10] we introduced the notion of theme which corresponds in the $[\lambda]$ -primitive case to frescos having a unique Jordan-Hölder sequence. Themes correspond to asymptotic expansion of a given vanishing period, so to the image of a fresco in the module of asymptotic expansions. For a fixed relative de Rham cohomology class (for instance given by a smooth differential form d -closed and df -closed) each choice of a vanishing cycle in the spectral eigenspace of the monodromy for the eigenvalue $\exp(-2i\pi.\lambda)$ produces a $[\lambda]$ -primitive theme, which is a quotient of the fresco associated to the given relative de Rham class itself. So the problem to determine which theme is a quotient of a given fresco is important to deduce possible asymptotic expansions of the various vanishing period integrals associated to a given relative de Rham class when we change the choice of the vanishing cycle.

In the appendix we prove a general existence result which naturally associate a fresco to any relative de Rham cohomology class of a proper holomorphic function of a complex manifold onto a disc.

AMS CLASSIFICATION. 32 S 25, 32 S 40, 32 S 50.

KEY WORDS. Theme, fresco, (a,b)-module, asymptotic expansion, vanishing period.

*Barlet Daniel, Institut Elie Cartan UMR 7502
Nancy-Université, CNRS, INRIA et Institut Universitaire de France,
BP 239 - F - 54506 Vandoeuvre-lès-Nancy Cedex.France. r
e-mail : barlet@iecn.u-nancy.fr

Contents

1	Introduction	2
2	Some known facts.	3
2.1	Regular and geometric (a,b)-modules.	3
2.2	Frescos and themes.	7
3	Commutation in Jordan-Hölder sequences.	8
3.1	The principal Jordan-Hölder sequence.	8
3.2	Commuting in $\tilde{\mathcal{A}}$	10
3.3	Some examples.	12
4	Semi-simple frescos.	15
4.1	The semi-simple filtration.	15
4.2	Co-semi-simple filtration.	21
4.3	Computation of the ss-depth.	23
4.4	Embedding for a semi-simple fresco.	24
5	Quotient themes of a $[\lambda]$-primitive fresco.	26
5.1	Structure theorem for $[\lambda]$ -primitive frescos.	26
5.2	Embedding dimension for a $[\lambda]$ -primitive fresco.	28
5.3	Quotient themes of a $[\lambda]$ -primitive fresco.	28
6	Appendix : the existence theorem.	32
6.1	Preliminaries.	32
6.2	$\tilde{\mathcal{A}}$ -structures.	35
6.3	The finiteness theorem.	38

1 Introduction

Let $f : X \rightarrow D$ be an holomorphic function on a connected complex manifold. Assume that $\{df = 0\} \subset \{f = 0\} := X_0$. We consider X as a degenerating family of complex manifolds parametrized by $D^* := D \setminus \{0\}$ with a singular member X_0 at the origin of D . Let ω be a smooth $(p+1)$ -differential form on X satisfying $d\omega = 0 = df \wedge \omega$. Then in many interesting cases (see for instance [JAG.08] , [B.III] for the case of a function with 1-dimensional singular set and the Appendix for the proper case) the relative family of de Rham cohomology classes induced on the fibers $X_s, s \in D^*$ of f by ω/df is solution of a minimal filtered differential equation defined from the Gauss-Manin connexion of f . This object, called a **fresco** is a monogenic regular (a,b)-module satisfying an extra condition, called "geometric", which encodes simultaneously the regularity at 0 of the Gauss-Manin connexion, the monodromy theorem and B. Malgrange's positivity theorem.

We study the structure of such an object in order to determine the possible quotient themes of a given fresco. Such a theme corresponds to a possible asymptotic expansion of vanishing periods constructed from ω by choosing a vanishing cycle $\gamma \in H_p(X_{s_0}, \mathbb{C})$ and putting

$$F_\gamma(s) := \int_{\gamma_s} \omega/df$$

where γ_s is the (multivalued) horizontal family of cycles defined from γ in the fibers of f (see [M.74]).

We obtain a rather precise description of the themes of these vanishing periods in term of the structure of the fresco associated to ω .

We give in the Appendix the existence theorem of the fresco associated to a smooth d -closed and df -closed form in the case of a proper holomorphic function on a complex manifold. This key result was written in the preprint [B.08] and was not yet published.

It is also interesting to indicate to the reader that he may find some "algebraic" explicit computations in [B.09] of the fresco generated some monomial in the Brieskorn module of the isolated singularity $x^5 + y^5 + x^2.y^2$ which is one of the simplest example with a not semi-simple monodromy.

2 Some known facts.

2.1 Regular and geometric (a,b)-modules.

The main purpose of this paper is to give a precise description of the structure of a $[\lambda]$ -primitive monogenic and geometric (a,b)-module ; we shall call such an object a $[\lambda]$ -primitive **fresco**. It corresponds to the $[\lambda]$ -primitive part of the minimal filtered differential equation satisfied by a relative de Rham cohomology class as indicated in the introduction.

Let us first recall the definition of an (a,b)-module¹.

Définition 2.1.1 *An (a,b)-module E is a free finite rank $\mathbb{C}[[b]]$ -module endowed with an $\mathbb{C}$ -linear map $a : E \rightarrow E$ which satisfies the following two conditions :*

- *The commutation relation $a.b - b.a = b^2$.*
- *The map a is continuous for the b -adic topology of E .*

Remark that these two conditions imply that for any $S \in \mathbb{C}[[b]]$ we have

$$a.S(b) = S(b).a + b^2.S'(b)$$

¹For more details on these basic facts, see [B.93]

where S' is defined via the usual derivation on $\mathbb{C}[[b]]$. For a given free rank k $\mathbb{C}[[b]]$ -module with basis $e_1, \dots, e_k$, to define a structure of (a, b) -module it is enough to prescribe (arbitrarily) the values of a on $e_1, \dots, e_k$.

An alternative way to define (a, b) -modules is to consider the $\mathbb{C}$ -algebra

$$\tilde{\mathcal{A}} := \left\{ \sum_{\nu=0}^{\infty} P_{\nu}(a) \cdot b^{\nu} \right\}$$

where the P_{ν} are polynomials in $\mathbb{C}[z]$ and where the product by a is left and right continuous for the b -adic filtration and satisfies the commutation relation $a \cdot b - b \cdot a = b^2$.

Then a left $\tilde{\mathcal{A}}$ -module which is free and finite rank on the subalgebra $\mathbb{C}[[b]] \subset \tilde{\mathcal{A}}$ is an (a, b) -module and conversely.

An (a, b) -module E has a **simple pole** when we have $a \cdot E \subset b \cdot E$ and it is **regular** when it is contained in a simple pole (a, b) -module. The regularity is equivalent to the finiteness on $\mathbb{C}[[b]]$ of the saturation $E^{\#}$ of E by $b^{-1} \cdot a$ in $E \otimes_{\mathbb{C}[[b]]} \mathbb{C}[[b]][b^{-1}]$.

- Submodules and quotients of regular (a, b) -modules are regular.

Another important property of regular (a, b) -module is the existence of Jordan-Hölder sequences (J-H. sequences for short).

Recall first that any regular rank 1 (a, b) -module is characterized up to isomorphism, by a complex number λ and the corresponding isomorphism class is represented by the (a, b) -module $E_{\lambda} := \mathbb{C}[[b]] \cdot e_{\lambda}$ where $a \cdot e_{\lambda} = \lambda \cdot b \cdot e_{\lambda}$, which is isomorphic to the $\tilde{\mathcal{A}}$ -module $\tilde{\mathcal{A}} / \tilde{\mathcal{A}} \cdot (a - \lambda \cdot b)$.

Recall also that a submodule F of the (a, b) -module E is called **normal** when $F \cap b \cdot E = b \cdot F$. Normality is a necessary and sufficient condition in order that the quotient E/F is again an (a, b) -module.

A **Jordan-Hölder sequence** for the rank k regular (a, b) -module E is a sequence of normal submodules² $\{0\} = F_0 \subset F_1 \subset \dots \subset F_{k-1} \subset F_k = E$ such that the quotients F_j / F_{j-1} for $j \in [1, k]$ are rank 1 (a, b) -modules. So, to each J-H. sequence of E , we may associate an ordered sequence of complex numbers $\lambda_1, \dots, \lambda_k$ such $F_j / F_{j-1} \simeq E_{\lambda_j}$ for each $j \in [1, k]$.

Existence of J-H. sequence for any regular (a, b) -module and also the following lemma are proved in [B.93].

Lemme 2.1.2 *Let E be a regular (a, b) -module of rank k . Up to a permutation, the set $\{\exp(-2i\pi \cdot \lambda_j), j \in [1, k]\}$ is independant of the choice of the J-H. sequence of E . Moreover, the sum $\sum_{j=1}^k \lambda_j$ is also independant of the choice of the J-H. sequence of E .*

²For $G \subset F \subset E$ submodules with F normal in E , the normality of G in F is equivalent to the normality of G in E .

The **Bernstein polynomial** of a regular (a,b) -module E of rank k is defined as the minimal polynomial of $-b^{-1}.a$ acting on the k -dimensional $\mathbb{C}$ -vector space $E^\# / b.E^\#$. Of course, when E is the b -completion of the Brieskorn module of a non constant germ $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ of holomorphic function with an isolated singularity, we find the "usual" (reduced) Bernstein polynomial of f (see for instance [K.76] or [Bj.93]).

We say that a regular (a,b) -module E is **geometric** when all roots of its Bernstein polynomial are negative rational numbers. This condition which correspond to M. Kashiwara theorem [K.76], encodes the monodromy theorem and the positivity theorem of B. Malgrange (see [M.75] or the appendix of [B.84]) extending the situation of (a,b) -modules deduced from the Gauss-Manin connection of an holomorphic function.

Recall that the tensor product of two (a,b) -modules E and F (see [B.I]) is defined as the $\mathbb{C}[[b]]$ -module $E \otimes_{\mathbb{C}[[b]]} F$ with the $\mathbb{C}$ -linear endomorphism defined by the rule $a.(x \otimes y) = (a.x) \otimes y + x \otimes (a.y)$. The tensor product by a fix (a,b) -module preserves short exact sequences of (a,b) -modules and $E_\lambda \otimes E_\mu \simeq E_{\lambda+\mu}$. So the tensor product of two regular (a,b) -modules is again regular.

Définition 2.1.3 *Let E be a regular (a,b) -module. The dual E^* of E is defined as the $\mathbb{C}[[b]]$ -module $\text{Hom}_{\mathbb{C}[[b]]}(E, E_0)$ with the $\mathbb{C}$ -linear map given by*

$$(a.\varphi)(x) = a.\varphi(x) - \varphi(a.x)$$

where $E_0 := \tilde{\mathcal{A}} / \tilde{\mathcal{A}}.a \simeq \mathbb{C}[[b]].e_0$ with $a.e_0 = 0$.

It is an easy exercise to see that a acts and satisfies the identity $a.b - b.a = b^2$ on E^* with the previous definition. We have $E_\lambda^* \simeq E_{-\lambda}$ and the duality transforms a short exact sequence of (a,b) -modules in a short exact sequence. So the dual of a regular (a,b) -module is again regular. But the dual of a geometric (a,b) -module is almost never geometric. To use duality in the geometric case we shall combine it with tensor product with E_N where N is a big enough rational number. Then $E^* \otimes E_N$ is geometric and $(E^* \otimes E_N)^* \otimes E_N \simeq E$. We shall refer to this process as "twisted duality".

Define now the left $\tilde{\mathcal{A}}$ -module of "formal multivalued expansions"

$$\Xi := \bigoplus_{\lambda \in \mathbb{Q} \cap]0,1]} \Xi_\lambda \quad \text{with} \quad \Xi_\lambda := \bigoplus_{j \in \mathbb{N}} \mathbb{C}[[b]].s^{\lambda-1} \cdot \frac{(\text{Log } s)^j}{j!}$$

with the action of a given by

$$a.(s^{\lambda-1} \cdot \frac{(\text{Log } s)^j}{j!}) = \lambda \cdot [b.s^{\lambda-1} \cdot \frac{(\text{Log } s)^j}{j!} + b.s^{\lambda-1} \cdot \frac{(\text{Log } s)^{j-1}}{(j-1)!}]$$

for $j \geq 1$ and $a.s^{\lambda-1} = \lambda.b(s^{\lambda-1})$, with, of course, the commutation relations $a.S(b) = S(b).a + b^2.S'(b)$ for $S \in \mathbb{C}[[b]]$.

For any geometric (a,b)-module of rank k , the vector space $\text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi)$ is of dimension k and this functor transforms short exact sequences of geometric (a,b)-modules in short exact sequences of finite dimensional vector spaces (see [B.05] for a proof).

In the case of the Brieskorn module of an isolated singularity germ of an holomorphic function f at the origin of $\mathbb{C}^{n+1}$ this vector space may be identified with the n -th homology group (with complex coefficients) of the Milnor's fiber of f (see [B.05]). The correspondance is given by associating to a (vanishing) cycle γ the $\tilde{\mathcal{A}}$ -linear map

$$[\omega] \mapsto \left[\int_{\gamma_s} \omega / df \right] \in \Xi$$

where $\omega \in \Omega_0^{n+1}$, γ_s is the multivalued horizontal family of n -cycles defined by γ in the fibers of f , and $[g]$ denotes the formal asymptotic expansion at $s = 0$ of the multivalued holomorphic function g .

Définition 2.1.4 *A regular (a,b)-module is $[\lambda]$ -primitive (resp. $[\Lambda]$ -primitive), where $[\lambda]$ is an element (resp. a subset) in $\mathbb{C}/\mathbb{Z}$, if all roots of its Bernstein polynomial are in $[-\lambda]$ (resp. in $[-\Lambda]$).*

If we have a short exact sequence of (a,b)-modules

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

with E regular (resp. geometric, resp. $[\lambda]$ -primitive) then F and G are regular (resp. geometric, resp. $[\lambda]$ -primitive).

Conversely if F and G are regular (resp. geometric, resp. $[\lambda]$ -primitive) then E is regular (resp. geometric, resp. $[\lambda]$ -primitive).

This implies that E is $[\lambda]$ -primitive if and only if it admits a J-H. sequence such that all numbers $\lambda_1, \dots, \lambda_k$ are in $[\lambda]$. And then any J-H. sequence of E has this property.

The following proposition is proved in [B.09] section 1.3.

Proposition 2.1.5 *Let E be a regular (a,b)-module and fix a subset Λ in $\mathbb{C}/\mathbb{Z}$. Then there exists a maximal submodule $E[\Lambda]$ in E which is $[\Lambda]$ -primitive. This submodule is normal in E .*

If $\{[\lambda_1], \dots, [\lambda_d]\}$ is the image in $\mathbb{C}/\mathbb{Z}$ of the set of the opposite of roots of the Bernstein polynomial of E , given with an arbitrary order, there exists a unique sequence $0 = F_0 \subset F_1 \subset F_2 \subset F_d = E$ of normal submodules of E such that F_j/F_{j-1} is $[\lambda_j]$ -primitive for each $j \in [1, d]$.

We call $E[\Lambda]$ the $[\Lambda]$ -**primitive part** of E .

Thanks to this result, to understand what are the possible $[\lambda]$ –primitive themes which are quotient of a given fresco, it will be enough to work with $[\lambda]$ –primitive frescos.

Remark also that, in the geometric situation, the choice of a vanishing cycle which belongs to the generalized eigenspace of the monodromy for the eigenvalue $\exp(-2i\pi.\lambda)$ produces vanishing periods with $[\lambda]$ –primitive themes.

2.2 Frescos and themes.

Here we recall some results from [B.09] and [B.10].

Définition 2.2.1 *We shall call **fresco** a geometric (a,b) -module which is generated by one element as an $\tilde{\mathcal{A}}$ –module.*

Définition 2.2.2 *We shall call a **theme** a fresco which is a sub- $\tilde{\mathcal{A}}$ –module of Ξ .*

Recall that a normal submodule and a quotient by a normal submodule of a fresco (resp. of a theme) is a fresco (resp. is a theme).

A regular rank 1 (a,b) -module is a fresco if and only if it is isomorphic to E_λ for some $\lambda \in \mathbb{Q}^{+*}$. All rank 1 frescos are themes. The classification of rank 2 regular (a,b) -modules given in [B.93] gives the list of $[\lambda]$ –primitive rank 2 frescos which is the following, where $\lambda_1 > 1$ is a rational number :

$$E = E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - (\lambda_1 - 1).b) \quad (1)$$

$$E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(1 + \alpha.b^p)^{-1}.(a - (\lambda_1 + p - 1).b) \quad (2)$$

where $p \in \mathbb{N} \setminus \{0\}$ and $\alpha \in \mathbb{C}$.

The themes in this list are these in (1) and these in (2) with $\alpha \neq 0$. For a $[\lambda]$ –primitive theme in case (2) the number $\alpha \neq 0$ will be called the **parameter** of the theme.

For frescos we have a more precise result on the numbers associated to a J-H. sequence :

Proposition 2.2.3 *Let E be a $[\lambda]$ –primitive fresco and $\lambda_1, \dots, \lambda_k$ be the numbers associated to a J-H. sequence of E . Then, up to a permutation, the numbers $\lambda_j + j, j \in [1, k]$ are independant of the choice of the J-H. sequence.*

The following structure theorem for frescos will be usefull in the sequel.

Théorème 2.2.4 (see [B.09] th.3.4.1) *Let E be a fresco of rank k . Then there exists an element $P \in \tilde{\mathcal{A}}$ which may be written as*

$$P := (a - \lambda_1.b)S_1^{-1}.(a - \lambda_2.b) \dots S_{k-1}^{-1}.(a - \lambda_k.b)$$

such that E is isomorphic to $\tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$. Here $S_1, \dots, S_{k-1}$ are elements in $\mathbb{C}[b]$ such that $S_j(0) = 1$ for each $j \in [1, k-1]$. The element $P_E := (a - \lambda_1.b) \dots (a - \lambda_k.b)$ of $\tilde{\mathcal{A}}$ is homogeneous in (a, b) and gives the Bernstein polynomial B_E de E via the formula

$$(-b)^{-k}.P_E = B_E(-b^{-1}.a).$$

So $P_E \in \tilde{\mathcal{A}}$ depends only on the isomorphism class of E .

Note that in the case of a fresco the Bernstein polynomial of E is equal to the **characteristic polynomial** of the action of $-b^{-1}.a$ on $E^\sharp$. This allows a nice formula for a short exact sequence of frescos :

Proposition 2.2.5 (see [B.09] prop.3.4.4) *Let $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ be a short exact sequence of frescos. Then we have the equality in $\tilde{\mathcal{A}}$:*

$$P_E = P_F.P_G \quad \text{which is equivalent to} \quad B_E(x) = B_F(x - rk(G)).B_G(x).$$

The situation for a $[\lambda]$ -primitive theme is more rigid :

Proposition 2.2.6 *A fresco E is a $[\lambda]$ -primitive theme if and only if it admits a unique normal rank 1 submodule. In this case the J-H. sequence is unique and contains all normal submodules of E . The corresponding numbers $\lambda_1, \dots, \lambda_k$ are such that the sequence $\lambda_j + j$ is increasing (may-be not strictly).*

3 Commutation in Jordan-Hölder sequences.

In this section we shall study the possible different J-H. sequences of a given $[\lambda]$ -primitive fresco. Thanks to proposition 2.1.5, it is easy to see that the $[\lambda]$ -primitive assumption does not reduce the generality of this study.

3.1 The principal Jordan-Hölder sequence.

Définition 3.1.1 *Let E be a $[\lambda]$ -primitive fresco of rank k and let*

$$0 = F_0 \subset F_1 \subset \dots \subset F_k = E$$

*be a J-H. sequence of E . Then for each $j \in [1, k]$ we have $F_j/F_{j-1} \simeq E_{\lambda_j}$, where $\lambda_1, \dots, \lambda_k$ are in $\lambda + \mathbb{N}$. We shall say that such a J-H. sequence is **principal** when the sequence $[1, k] \ni j \mapsto \lambda_j + j$ is increasing.*

It is proved in [B.09] prop. 3.5.2 that such a principal J-H. sequence exists for any $[\lambda]$ -primitive fresco. Moreover, the corresponding sequence $\lambda_1, \dots, \lambda_k$ is unique. The following proposition shows much more.

Proposition 3.1.2 *Let E be a $[\lambda]$ -primitive fresco. Then its principal J-H. sequence is unique.*

We shall prove the uniqueness by induction on the rank k of E .
We begin by the case of rank 2.

Lemme 3.1.3 *Let E be a rank 2 $[\lambda]$ -primitive fresco and let λ_1, λ_2 the numbers corresponding to a principal J-H. sequence of E (so $\lambda_1 + 1 \leq \lambda_2 + 2$). Then the normal rank 1 submodule of E isomorphic to E_{λ_1} is unique.*

PROOF. The case $\lambda_1 + 1 = \lambda_2 + 2$ is obvious because then E is a $[\lambda]$ -primitive theme (see [B.10] corollary 2.1.7). So we may assume that $\lambda_2 = \lambda_1 + p_1 - 1$ with $p_1 \geq 1$ and that E is the quotient $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - \lambda_2.b)$ (see the classification of rank 2 frescos in 2.2), because the result is clear when E is a theme. We shall use the $\mathbb{C}[[b]]$ -basis e_1, e_2 of E where a is defined by the relations

$$(a - \lambda_2.b).e_2 = e_1 \quad (a - \lambda_1.b).e_1 = 0.$$

This basis comes from the isomorphism above $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_1.b).(a - \lambda_2.b)$ deduced from the classification of rank 2 frescos with $e_2 = [1]$ and $e_1 = (a - \lambda_2.b).e_2$. Let look for $x := U.e_2 + V.e_1$ such that $(a - \lambda_1.b).x = 0$. Then we obtain

$$b^2.U'.e_2 + U.(a - \lambda_2.b).e_2 + (\lambda_2 - \lambda_1).b.U.e_2 + b^2.V'.e_1 = 0$$

which is equivalent to the two equations :

$$b^2.U' + (p_1 - 1).b.U = 0 \quad \text{and} \quad U + b^2.V' = 0.$$

The first equation gives $U = 0$ for $p_1 \geq 2$ and $U \in \mathbb{C}$ for $p_1 = 1$. As the second equation implies $U(0) = 0$, in all cases $U = 0$ and $V \in \mathbb{C}$. So the solutions are in $\mathbb{C}.e_1$. ■

Remark that in the previous lemma, if we assume $p_1 \geq 1$ and E is not a theme, it may exist infinitely many different normal (rank 1) submodules isomorphic to E_{λ_2+1} . But then, $\lambda_2 + 2 > \lambda_1 + 1$. See remark 2 following 3.2.1

PROOF OF PROPOSITION 3.1.2. As the result is obvious for $k = 1$, we may assume $k \geq 2$ and the result proved in rank $\leq k - 1$. Let $F_j, j \in [1, k]$ and $G_j, j \in [1, k]$ two J-H. principal sequences for E . As the sequences $\lambda_j + j$ and $\mu_j + j$ coincide up to the order and are both increasing, they coincide. Now let j_0 be the first integer in $[1, k]$ such that $F_{j_0} \neq G_{j_0}$. If $j_0 \geq 2$ applying the induction hypothesis to E/F_{j_0-1} gives $F_{j_0}/F_{j_0-1} = G_{j_0}/F_{j_0-1}$ and so $F_{j_0} = G_{j_0}$. So we may assume that $j_0 = 1$. Let H be the normalization³ of $F_1 + G_1$. As F_1

³The smallest normal submodule containing $F_1 + G_1$; it has the same rank than $F_1 + G_1$.

and G_1 are normal rank 1 and distinct, then H is a rank 2 normal submodule. It is a $[\lambda]$ -primitive fresco of rank 2 with two normal rank 1 sub-modules which are isomorphic as $\lambda_1 = \mu_1$. Moreover the principal J-H. sequence of H begins by a normal submodule isomorphic to E_{λ_1} . So the previous lemma implies $F_1 = G_1$. So for any $j \in [1, k]$ we have $F_j = G_j$. $\blacksquare$

Définition 3.1.4 *Let E be a $[\lambda]$ -primitive fresco and consider a J-H. sequence $F_j, j \in [1, k]$ of E . Put $F_j/F_{j-1} \simeq E_{\lambda_j}$ for $j \in [1, k]$ (with $F_0 = \{0\}$). We shall call **fundamental invariants** of E the (unordered) k -tuple $\{\lambda_j + j, j \in [1, k]\}$.*

Of course this definition makes sens because we know that this (unordered) k -tuple is independant of the choice of the J-H. of E .

Note that when E is a theme, the uniqueness of the J-H. gives a natural order on this k -tuple. So in the case of a theme the fundamental invariants will be an ordered k -tuple. With this convention, this is compatible with the definition of the fundamental invariants of a $[\lambda]$ -primitive theme given in [B.10], up to a shift.

In the opposite direction, when E is a semi-simple $[\lambda]$ -primitive fresco we shall see (in section 4) that any order of this k -tuple may be realized by a J-H. sequence of E .

3.2 Commuting in $\tilde{\mathcal{A}}$.

Let E be a $[\lambda]$ -primitive fresco. Any isomorphism $E \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ where $P \in \tilde{\mathcal{A}}$ is given by

$$P := (a - \lambda_1.b).S_1^{-1}.(a - \lambda_2.b) \dots S_{k-1}^{-1}.(a - \lambda_k.b)$$

determines a J-H. sequence for E associated to the $\mathbb{C}[[b]]$ -basis $e_1, \dots, e_k$ such that the relations $(a - \lambda_j.b).e_j = S_{j-1}.e_{j-1}$ hold for $j \in [1, k]$ with the convention $e_0 = 0$, and the fact that e_k corresponds, via the prescribed isomorphism, to the class of 1 modulo $\tilde{\mathcal{A}}.P$.

Lemme 3.2.1 *Let p_1 and p_2 be two positive integers and let $\lambda_1 \in \lambda + 2 + \mathbb{N}$, where $\lambda \in]0, 1] \cap \mathbb{Q}$. Define $P \in \tilde{\mathcal{A}}$ as*

$$P := (a - \lambda_1.b)S_1^{-1}(a - \lambda_2.b)S_2^{-1}(a - \lambda_3.b)$$

where $\lambda_{j+1} := \lambda_j + p_j - 1$ for $j = 1, 2$, and where S_1, S_2 lie in $\mathbb{C}[[b]]$ and satisfy $S_1(0) = S_2(0) = 1$. We assume that the coefficient of b^{p_1} in S_1 vanishes and that the coefficient of b^{p_2} of S_2 is $\alpha \neq 0$.

Then if $U \in \mathbb{C}[[b]]$ is any solution of the differential equation $b.U' = p_1.(U - S_1)$, we have

$$P = U^{-1}.(a - (\lambda_2 + 1).b).(S_1.U^{-2})^{-1}.(a - (\lambda_1 - 1).b).(U.S_2)^{-1}.(a - \lambda_3.b).$$

Moreover, there exists a unique choice of U such that the coefficient of $b^{p_1+p_2}$ in $U.S_2$ vanishes.

PROOF. The fact that a solution U of this differential equation satisfies

$$(a - \lambda_1.b)S_1^{-1}(a - \lambda_2.b) = U^{-1}.(a - (\lambda_2 + 1).b).S_1^{-1}.U^2.(a - (\lambda_1 - 1).b).U^{-1}$$

is proved in [B.09] lemma 3.5.1. We use here the case $\delta := \lambda - \mu = \lambda_2 + 1 - \lambda_1 = p_1$ and the fact that the coefficient of b^δ in S_1 vanishes.

As the solution U is unique up to $\mathbb{C}.b^{p_1}$, to prove the second assertion let U_0 be the solution with no term in b^{p_1} . Now the coefficient $\beta(\rho)$ of $b^{p_1+p_2}$ in $S_2.U$ where $U := U_0 + \rho.b^{p_1}$, is $\beta(\rho) = \beta(0) + \rho.\alpha$. As we assumed that $\alpha \neq 0$ there exists an unique choice of ρ for which $\beta(\rho) = 0$. ■

REMARKS.

1. In the situation of the previous lemma the rank 3 fresco $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ is an extension

$$0 \rightarrow E_{\lambda_1} \rightarrow E \rightarrow T_{\lambda_2, p_2}(\alpha) \rightarrow 0$$

where $T_{\lambda_2, p_2}(\alpha)$ is the rank 2 theme with fundamental invariants (λ_2, p_2) and parameter α (see the definition 2.2.2), so

$$T_{\lambda_2, p_2}(\alpha) \simeq \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.(a - \lambda_2.b).(1 + \alpha.b^{p_2})^{-1}.(a - (\lambda_2 + p_2 - 1).b).$$

2. Let ξ be in $\mathbb{C}^*$ and choose $\rho := (\xi - \beta_0)/\alpha$ in the previous proof. Then E is a extension

$$0 \rightarrow E_{\lambda_2+1} \rightarrow E \rightarrow T_{\lambda_1-1, p_1+p_2}(\xi) \rightarrow 0$$

where $T_{\lambda_1-1, p_1+p_2}(\xi)$ is the rank 2 theme with fundamental invariants $(\lambda_1 - 1, p_1 + p_2)$ and parameter ξ . This shows that we may have infinitely many non isomorphic rank 2 themes as quotients of a given rank 3 $[\lambda]$ -primitive fresco E . We have also infinitely many different J-H. sequences with the same quotients: $(\lambda_2 + 1, \lambda_1 - 1, \lambda_3)$.

Note that in this situation we have⁴ $\dim_{\mathbb{C}} [Ker(a - (\lambda_2 + 1).b)] = 2$.

Corollaire 3.2.2 *In the situation of the previous lemma choose $\rho = -\beta(0).\alpha$ and denote by V a solution⁵ of the differential equation $b.V' = (p_1 + p_2).(V - U.S_2)$. Then P is equal to*

$$U^{-1}.(a - (\lambda_2 + 1).b).S_1^{-1}.U^2.V^{-1}.(a - (\lambda_3 + 1).b).(US_2V^{-2})^{-1}.(a - (\lambda_1 - 2).b).V^{-1}. \quad (@)$$

Moreover, the coefficient of b^{p_2} in $S_1.U^{-2}.V$ is $(p_1 + p_2).\alpha/p_1$.

⁴It is easy to see that if Z is a solution in $\mathbb{C}[[b]]$ of the differential equation $b.Z' - p_1.Z + S_1 = 0$, then

$$Ker(a - (\lambda_2 + 1).b) = \{(r, s) \in \mathbb{C}^2 / r.(Z.e_1 + b.e_2) + s.b^{p_1}.e_1\}.$$

⁵Note that as $U.S_2$ has no term in $b^{p_1+p_2}$ with our choice of ρ , such a solution exists in $\mathbb{C}[[b]]$. Moreover $V(0) = (U.S_2)(0) = 1$ because $U(0) = S_1(0) = 1 = S_2(0)$.

PROOF. Of course the choice of ρ allows to apply again the lemma 3.5.1. of [B.09], with now $\delta = \lambda_3 + 1 - (\lambda_1 - 1) = p_1 + p_2$. This gives (@). Using $b.U' = p_1.(U - S_1)$ we get

$$\begin{aligned} b.U'.U^{-2} &= p_1.(U^{-1} - S_1.U^{-2}) \quad \text{and with } Z := U^{-1} \\ b.Z' &= -p_1.(Z - S_1.U^{-2}) \quad \text{and then} \\ b.Z'.V &= -p_1.(Z.V - S_1.U^{-2}.V) \end{aligned} \quad (@@)$$

But using also $b.V' = (p_1 + p_2).(V - U.S_2)$ we get

$$b.V'.Z = (p_1 + p_2).(V.Z - S_2).$$

Adding with (@@) gives

$$b.(V.Z)' - p_2.V.Z = p_1.S_1.U^{-2}.V - (p_1 + p_2).S_2$$

which leads to the result, because the left handside has no term in b^{p_2} . ■

An obvious consequence of this corollary is that there exists in E a normal subtheme isomorphic to $T_{\lambda_2+1,p_2}((1 + p_2/p_1).\alpha)$, so with fundamental invariants $(\lambda_2 + 1, p_2)$ and with parameter $(1 + p_2/p_1).\alpha$. Recall that $T_{\lambda_2,p_2}(\alpha)$ was the rank 2 quotient theme which appears in the principal J-H. of E .

3.3 Some examples.

We shall give here some examples, showing the complexity of the non commutative structure of the algebra $\tilde{\mathcal{A}}$.

Lemme 3.3.1 *Let x, y, z non zero complex numbers, λ_1 a rational number bigger than 3 and p_1, p_2, p_3 three positive distinct integers. Assume that p_3 is not a multiple of p_2 , and define $\lambda_{j+1} := \lambda_j + p_j - 1$ for $j = 1, 2, 3$. Put*

$$R_1 := 1 + x.b^{p_1}, \quad R_3 := 1 + y.b^{p_3} + z.b^{p_2+p_3}, \quad U := 1 - \frac{z}{y}.b^{p_2},$$

$$S := R_1.U \quad \text{and} \quad T := U.R_3.$$

Then T has no term in $b^{p_2+p_3}$ and there exists a solution $V \in \mathbb{C}[[b]]$ of the differential equation $b.V' = (p_2 + p_3).(V - T)$.

Then the element $P := (a - \lambda_1.b).R_1^{-1}.(a - \lambda_2.b).(a - \lambda_3.b).R_3^{-1}.(a - \lambda_4.b)$ in $\tilde{\mathcal{A}}$ is equal to

$$(a - \lambda_1.b).S^{-1}.(a - (\lambda_3 + 1).b).U^2.V^{-1}.(a - (\lambda_4 + 1).b).T^{-1}.V^2.(a - (\lambda_2 - 2).b).V^{-1}. \quad (1)$$

PROOF. A simple computation gives

$$T = 1 - \frac{z}{y}.b^{p_2} + y.b^{p_3} - \frac{z^2}{y}.b^{2p_2+p_3}$$

and

$$V = 1 - \frac{z}{y} \frac{p_2 + p_3}{p_3}.b^{p_2} + \frac{p_2 + p_3}{p_2}.y.b^{p_3} + \rho.b^{p_2+p_3} + p_2 \frac{z^2}{y}.b^{2p_2+p_3}$$

where ρ is an arbitrary complex number.

Using the lemma 3.5.1. of [B.09] and the fact that U satisfies $b.U' = p_2.(U - 1)$ we get

$$P = (a - \lambda_1.b).S^{-1}.(a - (\lambda_3 + 1).b).U^2.(a - (\lambda_2 - 1).b).T^{-1}.(a - \lambda_4.b).$$

As $\lambda_4 = (\lambda_2 - 1) + p_2 + p_3 - 1$ and T has no term in $b^{p_2+p_3}$, we obtain, using again the lemma of *loc. cit.*

$$(a - (\lambda_2 - 1).b).T^{-1}.(a - \lambda_4.b) = V^{-1}.(a - (\lambda_4 + 1).b)T^{-1}.V^2.(a - (\lambda_2 - 2).b).V^{-1}$$

if V is a solution of $b.V' = (p_2 + p_3).(V - T)$; this implies (1). ■

Lemme 3.3.2 *In the situation of the previous lemma the rank 4 fresco given by $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ is not a theme, but we have the following exact sequences :*

$$\begin{aligned} 0 \rightarrow T_1 \rightarrow E \rightarrow T_2 \rightarrow 0 \\ 0 \rightarrow T_3 \rightarrow E \rightarrow E_{\lambda_2-2} \rightarrow 0. \end{aligned}$$

where T_1 and T_2 are rank 2 themes and T_3 a rank 3 theme.

PROOF. The first exact sequence is consequence of the definition of P , and the rank 2 theme T_1 has (λ_1, p_1) as fundamental invariants and x as parameter ; the rank 2 theme T_2 has (λ_3, p_3) as fundamental invariants and y as parameter. Let e be a generator of E whose annihilator is $\tilde{\mathcal{A}}.P$. Then the relation (1) shows that $\varepsilon := T^{-1}.V^2.(a - (\lambda_2 - 2).b).V^{-1}.e$ in E is annihilated by

$$Q := (a - \lambda_1.b).S^{-1}.(a - (\lambda_3 + 1).b).U^2.V^{-1}.(a - (\lambda_4 + 1).b).$$

So $\tilde{\mathcal{A}}.\varepsilon$ has rank 3 and is normal because $E/\tilde{\mathcal{A}}.\varepsilon \simeq E_{\lambda_2-2}$. We shall prove that $\tilde{\mathcal{A}}.\varepsilon$ is a theme. As $\lambda_3 + 1 = \lambda_1 + p_1 + p_2 - 1$ and $\lambda_4 + 1 = (\lambda_3 + 1) + p_3 - 1$, it is enough to check that the coefficient of $b^{p_1+p_2}$ in S and the coefficient in b^{p_3} in $U^{-2}.V$ do not vanish. As we have

$$\text{i) } S = 1 + x.b^{p_1} - \frac{z}{y}.b^{p_2} - \frac{xz}{y}.b^{p_1+p_2}$$

$$\text{ii) } V = 1 - \frac{z}{y} \frac{p_2+p_3}{p_3}.b^{p_2} + \frac{p_2+p_3}{p_2}.y.b^{p_3} + \rho.b^{p_2+p_3} + p_2 \frac{z^2}{y}.b^{2p_2+p_3}$$

$$\text{iii) } U^{-2} = \sum_{n=1}^{\infty} n. \left(\frac{z}{y} b^{p_2} \right)^{n-1}$$

these coefficients are respectively equal to $-\frac{x.z}{y}$ and $\frac{p_2+p_3}{p_2}.y$ using the fact that p_3 is not a multiple of p_2 . ■

REMARK. Choosing for instance $U = 1$ gives

$$P = (a - \lambda_1.b).R_1^{-1}.(a - (\lambda_3 + 1).b).(a - (\lambda_2 - 1).b).R_3^{-1}.(a - \lambda_4.b)$$

and then if W is a solution of the differential equation $b.W' = (p_1 + p_2).(W - R_1)$ we obtain

$$P = W^{-1}.(a - (\lambda_3 + 2).b).(R_1.W^{-2})^{-1}.(a - (\lambda_1 - 1).b).W^{-1}.(a - (\lambda_2 - 1).b).R_3^{-1}.(a - \lambda_4.b)$$

and we have an exact sequence

$$0 \rightarrow E_{\lambda_3+2} \rightarrow E \rightarrow T_4 \rightarrow 0$$

where T_4 is a rank 3 theme and where the corresponding J-H. sequence associated to this exact sequence satisfies $F_2 = S_1(E)$ where $S_1(E)$ is the maximal semi-simple normal submodule of E (see section 4) .

Note that the first exact sequence corresponds to the principal J-H. of E .

The second gives a J-H. sequence such that its quotients correspond to the order $\lambda_1 + 1, \lambda_3 + 3, \lambda_4 + 4, \lambda_2 + 2$ of the increasing sequence $\lambda_j + j, j \in [1, 4]$. The last sequence above corresponds to the order $\lambda_3 + 3, \lambda_1 + 1, \lambda_2 + 2, \lambda_4 + 4$. In this example the semi-simple depth $d(E)$ of E (see section 4) is equal to 3.

EXEMPLE. We give here an example a $[\lambda]$ -primitive fresco of rank 4 with a J-H. sequence having no non commuting index but which is not semi-simple (see the section 4 below).

Let $\lambda_1 > 4$ be a rational number and p_1, p_2, p_3 be strictly positive integers. Then consider the fresco

$$E := \tilde{\mathcal{A}} / (a - \lambda_1.b).(a - \lambda_2.b).(1 + b^{p_2+p_3})^{-1}.(a - \lambda_3.b).(a - \lambda_4.b)$$

where we define $\lambda_{j+1} = \lambda_j + p_j - 1$ for $j = 1, 2, 3$. Then it is clear that all indices of the principal J-H. sequence of E are commuting indices : for $i = 1$ and $i = 3$ this is obvious, for $i = 2$ this results from the commuting lemme 3.5.1 of [B.09] and the fact that $p_2 + p_3 > p_3$ as we assume $p_2 \geq 1$. Now we have the equality in $\tilde{\mathcal{A}}$:

$$\begin{aligned} (a - \lambda_1.b).(a - \lambda_2.b).(1 + b^{p_1+p_2})^{-1}.(a - \lambda_3.b).(a - \lambda_4.b) = \\ (a - \lambda_1.b).(a - \lambda_2.b).(1 + b^{p_1+p_2})^{-1}.(a - (\lambda_4 + 1).b).(a - (\lambda_3 - 1).b). \end{aligned}$$

This shows, because $\lambda_4 + 1 = \lambda_2 + p_2 + p_3 - 1$, that there exists a subquotient of rank 2 of E which is a theme; so E is not semi-simple. In fact, we produce another J-H. sequence with one non commuting index !

4 Semi-simple frescos.

4.1 The semi-simple filtration.

Définition 4.1.1 *We shall say that a fresco E is **semi-simple** if any quotient of E which is a $[\lambda]$ -primitive theme for some $[\lambda] \in \mathbb{Q}/\mathbb{Z}$ is of rank ≤ 1 .*

REMARKS.

1. A $[\lambda]$ -primitive theme is semi-simple if and only if it has rank ≤ 1 .
2. An equivalent definition of a semi-simple fresco is to ask that any $\tilde{\mathcal{A}}$ -linear map

$$\varphi : E \rightarrow \Xi_\lambda$$

for some $[\lambda] \in \mathbb{Q}/\mathbb{Z}$ has rank ≤ 1 . This is a necessary condition because $\varphi(E)$ is a $[\lambda]$ -primitive theme which is a quotient of E . The converse comes from the fact that any $[\lambda]$ -primitive theme admits an injective $\tilde{\mathcal{A}}$ -linear map in Ξ_λ .

3. A fresco is semi-simple if and only if for each $[\lambda]$ its $[\lambda]$ -primitive part (see the proposition 2.1.5) is semi-simple : if $E[\lambda]$ is the $[\lambda]$ -primitive part of E the restriction map $\text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda) \rightarrow \text{Hom}_{\tilde{\mathcal{A}}}(E[\lambda], \Xi_\lambda)$ is an isomorphism. For instance, a theme with only rank ≤ 1 $[\lambda]$ -primitive part for each $[\lambda] \in \mathbb{C}/\mathbb{Z}$ is semi-simple.

Lemme 4.1.2 *For any $F \subset E$ a normal submodule of a semi-simple fresco E , F and E/F are semi-simple frescos. So any sub-quotient⁶ and of a semi-simple fresco is again a semi-simple fresco.*

PROOF. As any $\tilde{\mathcal{A}}$ -linear map $\psi : F \rightarrow \Xi_\lambda$ extends to a $\tilde{\mathcal{A}}$ -linear map $\varphi : E \rightarrow \Xi_\lambda$ (see section 2 or [B.05]) the semi-simplicity of E implies the semi-simplicity of F . The semi-simplicity of E/F is obvious. ■

Corollaire 4.1.3 *Let E be a semi-simple fresco with rank k and let $\lambda_1, \dots, \lambda_k$ be the numbers associated to a J-H. sequence of E . Let $\mu_1, \dots, \mu_k$ be a twisted permutation⁷ of $\lambda_1, \dots, \lambda_k$. Then there exists a J-H. sequence for E with quotients corresponding to $\mu_1, \dots, \mu_k$.*

⁶By a subquotient H we mean that there exists $G \subset F$ normal submodules in E such that $H := F/G$. Remark that H is a quotient of a normal submodule but also a submodule of a quotient of E , as $F/G \subset E/G$.

⁷This means that the sequence $\mu_j + j, j \in [1, k]$ is a permutation (in the usual sens) of $\lambda_j + j, j \in [1, k]$.

PROOF. As the symmetric group $\mathfrak{S}_k$ is generated by the transpositions $t_{j,j+1}$ for $j \in [1, k-1]$, it is enough to show that, if E has a J-H. sequence with quotients given by the numbers $\lambda_1, \dots, \lambda_k$ then there exists a J-H. sequence for E with quotients $\lambda_1, \dots, \lambda_{j-1}, \lambda_{j+1} + 1, \lambda_j - 1, \lambda_{j+2}, \dots, \lambda_k$ for $j \in [1, k-1]$. But $G := F_{j+1}/F_{j-1}$ is a rank 2 sub-quotient of E with an exact sequence

$$0 \rightarrow E_{\lambda_j} \rightarrow G \rightarrow E_{\lambda_{j+1}} \rightarrow 0.$$

As G is a rank 2 semi-simple fresco, it admits also an exact sequence

$$0 \rightarrow G_1 \rightarrow G \rightarrow G/G_1 \rightarrow 0$$

with $G_1 \simeq E_{\lambda_{j+1}+1}$ and $G/G_1 \simeq E_{\lambda_j-1}$. Let $q : F_{j+1} \rightarrow G$ be the quotient map. Now the J-H. sequence for E given by

$$F_1, \dots, F_{j-1}, q^{-1}(G_1), F_{j+1}, \dots, F_k = E$$

satisfies our requirement. ■

Proposition 4.1.4 *Let E be a $[\lambda]$ -primitive fresco. A necessary and sufficient condition in order that E is semi-simple is that it admits a J-H. sequence with quotient corresponding to $\mu_1, \dots, \mu_k$ such that the sequence $\mu_j + j$ is strictly decreasing.*

REMARKS.

1. As a fresco is semi-simple if and only if for each $[\lambda]$ its $[\lambda]$ -primitive part is semi-simple, this proposition gives also a criterium to semi-simplicity for any fresco.
2. This criterium is a very efficient tool to produce easily examples of semi-simple frescos.

PROOF. Remark first that if we have, for a $[\lambda]$ -primitive fresco E , a J-H. sequence $F_j, j \in [1, k]$ such that $\lambda_j + j = \lambda_{j+1} + j + 1$ for some $j \in [1, k-1]$, then F_{j+1}/F_{j-1} is a sub-quotient of E which is a $[\lambda]$ -primitive theme of rank 2. So E is not semi-simple, thanks to the previous corollary. So when a $[\lambda]$ -primitive fresco E is semi-simple the principal J-H. sequence corresponds to a strictly increasing sequence $\lambda_j + j$. Now, thanks again to the previous corollary we may find a J-H. sequence for E corresponding to the strictly decreasing order for the sequence $\lambda_j + j$.

No let us prove the converse. We shall use the following lemma.

Lemme 4.1.5 *Let F be a rank k semi-simple $[\lambda]$ -primitive fresco and let $\lambda_j + j$ the strictly increasing sequence corresponding to its principal J-H. sequence. Let $\mu \in [\lambda]$ such that $0 < \mu + k + 1 < \lambda_1 + 1$. Then any fresco E in an exact sequence*

$$0 \rightarrow F \rightarrow E \rightarrow E_\mu \rightarrow 0$$

is semi-simple (and $[\lambda]$ -primitive).

PROOF. Assume that we have a rank 2 quotient $\varphi : E \rightarrow T$ where T is a $[\lambda]$ -primitive theme. Then $\text{Ker } \varphi \cap F$ is a normal submodule of F of rank $k-2$ or $k-3$. If $\text{Ker } \varphi \cap F$ is of rank $k-3$, the rank of $F/\text{Ker } \varphi \cap F$ is 2 and it injects in T via φ . So $F/\text{Ker } \varphi \cap F$ is a rank 2 $[\lambda]$ -primitive theme. As it is semi-simple, because F is semi-simple, we get a contradiction.

So the rank of $F/\text{Ker } \varphi \cap F$ is 1 and we have an exact sequence

$$0 \rightarrow F/\text{Ker } \varphi \cap F \rightarrow T \rightarrow E/F \rightarrow 0.$$

Put $F/\text{Ker } \varphi \cap F \simeq E_\lambda$. Because T is a $[\lambda]$ -primitive theme, we have the inequality $\lambda+1 \leq \mu+2$. But we know that $\lambda_1+1 \leq \lambda+k$ because $\lambda+k$ is in the set $\{\lambda_j+j, j \in [1, k]\}$ and λ_1+1 is the infimum of this set. So $\lambda_1+1 \leq \mu+k+1$ contradicting our assumption that $\mu+k+1 < \lambda_1+1$. ■

END OF PROOF OF THE PROPOSITION 4.1.4. Now we shall prove by induction on the rank of a $[\lambda]$ -primitive fresco E that if it admits a J-H. sequence corresponding to a strictly decreasing sequence $\mu_j + j$, it is semi-simple. As the result is obvious in rank 1, we may assume $k \geq 1$ and the result proved for k . So let E be a fresco of rank $k+1$ and let $F_j, j \in [1, k+1]$ a J-H. sequence for E corresponding to the strictly decreasing sequence $\mu_j + j, j \in [1, k+1]$. Put $F_j/F_{j-1} \simeq E_{\mu_j}$ for all $j \in [1, k+1]$, define $F := F_k$ and $\mu := \mu_{k+1}$; then the induction hypothesis gives that F is semi-simple and we may apply the previous lemma. ■

Proposition 4.1.6 *Let E be a fresco. There exists a unique maximal normal semi-simple submodule $S_1(E)$ in E . It contains any (normal) submodule of rank 1 contained in E . Moreover, if $S_1(E)$ is of rank 1, then E is a $[\lambda]$ -primitive theme.*

PROOF. For any λ and any non zero $\varphi \in \text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda)$ let $F_1(\varphi)$ be the rank 1 submodule of the $[\lambda]$ -primitive theme $\varphi(E)$. Now put

$$S_1(E) := \bigcap_{\lambda} \bigcap_{\varphi \in \text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda) \setminus \{0\}} [\varphi^{-1}(F_1(\varphi))].$$

Let us prove that $S_1(E)$ is a normal semi-simple submodule. Normality is obvious as it is an intersection of normal submodules. To prove semi-simplicity, let $\psi : S_1(E) \rightarrow \Xi_\lambda$ be a $\tilde{\mathcal{A}}$ -linear map. Using the surjectivity of the restriction $\varphi \in \text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda) \rightarrow \varphi|_{S_1(E)} \in \text{Hom}_{\tilde{\mathcal{A}}}(S_1(E), \Xi_\lambda)$, (see section 2.1 or [B.05]), we see immediately that ψ has rank ≤ 1 . So $S_1(E)$ is semi-simple.

Now consider a semi-simple normal submodule S in E . For any $\varphi \in \text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda)$ the restriction of φ to S has rank ≤ 1 . So $\varphi(S)$ is contained in the normal rank 1 submodule $F_1(\varphi)$ of the $[\lambda]$ -primitive theme $\varphi(E)$. So S is contained in $\varphi^{-1}(F_1(\varphi))$ for each φ . Then $S \subset S_1(E)$, and this proves the maximality of $S_1(E)$.

Consider now any rank 1 normal submodule F of E . As F is semi-simple and

normal in E , we have $F \subset S_1(E)$. If $S_1(E)$ is rank 1, there exists an unique rank 1 normal submodule in E . Then E is a $[\lambda]$ -primitive theme, thanks to [B.10] theorem 2.1.6. ■

The following interesting corollary is an obvious consequence of the previous proposition.

Corollaire 4.1.7 *Let E be a fresco and let $\lambda_1, \dots, \lambda_k$ be the numbers associated to any J-H. sequence of E . Let $\mu_1, \dots, \mu_d$ be the numbers associated to any J-H. sequence of $S_1(E)$. Then, for $j \in [1, k]$, there exists a rank 1 normal submodule of E isomorphic to E_{λ_j+j-1} if and only if there exists $i \in [1, d]$ such that we have $\lambda_j + j - 1 = \mu_i + i - 1$.*

Of course, this gives the list of all isomorphy classes of rank 1 normal submodules contained in E . So, using shifted duality, we get also the list of all isomorphy classes of rank 1 quotients of E .

Définition 4.1.8 *Let E be a fresco. Define inductively the increasing sequence $S_j(E), j \geq 0$ of normal submodules of E by putting $S_0(E) := \{0\}$ and for $j \geq 1$ $S_j(E)/S_{j-1}(E) := S_1(E/S_{j-1}(E))$. We shall call $S_j(E), j \geq 0$ the **semi-simple filtration** of E . We shall call **semi-simple-depth** of E (**ss-depth** for short) the first integer $d = d(E) \geq 0$ such that $E = S_d(E)$.*

EXAMPLE. In the example of lemma 3.2.1 let F_2 the second step of the principal J-H. sequence of E . Then $F_2 = S_1(E)$ is the maximal semi-simple normal submodule of E . This is a consequence of the fact that E is not semi-simple, F_2 admits a J-H. sequence with quotients $E_{\lambda_2+1}, E_{\lambda_1-1}$ with $\lambda_2 + 2 > \lambda_1 + 1$, so we may apply proposition 4.1.4.

Proposition 4.1.9 *Let E be a fresco. Then we have the following properties :*

- i) *Any $[\lambda]$ -primitive sub-theme T in E of rank j is contained in $S_j(E)$.*
- ii) *Any $[\lambda]$ -primitive quotient theme T of $S_j(E)$ has rank $\leq j$.*
- iii) *For any $j \in \mathbb{N}$ we have*

$$S_j(E) = \cap_{\lambda} \cap_{\varphi \in \text{Hom}_{\tilde{A}}(E, \Xi_{\lambda})} [\varphi^{-1}(F_j(\varphi))]$$

where $F_j(\varphi)$ is the normal submodule of rank j of the $[\lambda]$ -primitive theme $\varphi(E)$, with the convention that $F_j(\varphi) = \varphi(E)$ when the rank of φ is $\leq j$.

- iv) *The ss-depth of E is equal to d if and only if d is the maximal rank of a $[\lambda]$ -primitive quotient theme of E .*
- v) *The ss-depth of E is equal to d if and only if d is the maximal rank of a normal $[\lambda]$ -primitive sub-theme of E .*

REMARKS.

1. By definition of the ss-depth $d(E)$ of E the semi-simple filtration is strictly increasing for $j \in [0, d(E)]$.
2. Let E be a fresco and $N \in \mathbb{Z}$ such that $E \otimes E_N$ is geometric (so is again a fresco). Then E is semi-simple (resp. a theme) if and only if $E \otimes E_N$ is semi-simple (resp. a theme). Moreover, in this situation we have $d(E) = d(E \otimes E_N)$.
3. Let F be a submodule in a fresco E , and denote $\tilde{F}$ its normalization. Then $\tilde{F}$ is monogenic (being normal in a monogenic) and geometric. As there exists $N \in \mathbb{N}$ such that $b^N \cdot \tilde{F} \subset F$, $\tilde{F}$ is a theme for F a theme. The analog result is also true for a semi-simple F : if $\varphi : \tilde{F} \rightarrow \Xi_\lambda$ has rank ≥ 2 , as F has finite codimension in $\tilde{F}$, the restriction of φ to F has also rank ≥ 2 which contradicts the semi-simplicity of F .

So we have proved the following two assertions :

- If $T \subset E$ is a theme in a fresco E , its normalization is also a theme (of same rank than T).
- If $S \subset E$ is a semi-simple fresco in a fresco E , its normalization is also a semi-simple fresco.

PROOF OF PROPOSITION 4.1.9. Let us prove i) by induction on j . As the case $j = 1$ is obvious, let us assume that $j \geq 2$ and that the result is proved for $j - 1$. Let T a $[\lambda]$ -primitive theme in E , and let $F_{j-1}(T)$ be its normal submodule of rank $j - 1$ (equal to T if the rank of T is less than $j - 1$). Then by the induction hypothesis, we have $F_{j-1}(T) \subset S_{j-1}(E)$. Then we have a $\tilde{\mathcal{A}}$ -linear map $T/F_{j-1}(T) \rightarrow E/S_{j-1}(E)$. If the rank of T is at most j , then $T/F_{j-1}(T)$ has rank at most 1 and its image is in $S_1(E/S_{j-1}(E))$. So $T \subset S_j(E)$.

To prove ii) we also make an induction on j . The case $j = 1$ is obvious. So we may assume $j \geq 2$ and the result proved for $j - 1$. Let $\varphi : S_j(E) \rightarrow T$ a surjective map on a $[\lambda]$ -primitive theme T . By the inductive hypothesis we have $\varphi(S_{j-1}(E)) \subset F_{j-1}(T)$. So we have an induced surjective map

$$\tilde{\varphi} : S_j(E)/S_{j-1}(E) \rightarrow T/F_{j-1}(T).$$

As $S_j(E)/S_{j-1}(E)$ is semi-simple, the image of $\tilde{\varphi}$ has rank ≤ 1 . It shows that T has rank $\leq j$.

To prove iii) consider first a $\tilde{\mathcal{A}}$ -linear map $\varphi : E \rightarrow \Xi_\lambda$. As $\varphi(E)$ is a $[\lambda]$ -primitive theme, $\varphi(S_j(E))$ is a $[\lambda]$ -primitive theme quotient of $S_j(E)$. So its rank is $\leq j$ and we have $\varphi(S_j(E)) \subset F_j(\varphi)$.

Conversely, for any $\tilde{\mathcal{A}}$ -linear map $\varphi : E \rightarrow \Xi_\lambda$, the image $\varphi(S_j(E))$ is a $[\lambda]$ -primitive quotient theme of $S_j(E)$. So its rank is $\leq j$ and it is contained in $F_j(\varphi)$.

Let us prove iv). If $S_d(E) = E$ then any $[\lambda]$ -primitive sub-theme in E has rank $\leq d$ thanks to ii). Conversely, assume that for any $[\lambda]$ any $[\lambda]$ -primitive sub-theme of E has rank $\leq d-1$ and $S_{d-1}(E) \neq E$. Then choose a $\tilde{\mathcal{A}}$ -linear map $\varphi : E \rightarrow \Xi_\lambda$ such that $\varphi^{-1}(F_{d-1}(\varphi)) \neq E$. Then $\varphi(E)$ is a $[\lambda]$ -primitive theme of rank d which is a quotient of E , thanks to the following lemma 4.1.10. To prove v) let us show that if E is a fresco and $N \gg 1$ an integer, then $E^* \otimes E_N$ is again a fresco and that we have the inequality $d(E^* \otimes E_N) \geq d(E)$. The fact that for N a large enough integer $E^* \otimes E_N$ is again a fresco is clear. Now, as E has a $[\lambda]$ -primitive quotient theme of rank d , then $E^* \otimes E_N$ has a $[-\lambda]$ -primitive sub-theme of rank d . So we obtain the inequality $d(E^* \otimes E_N) \geq d(E)$ from i). Now, using again duality and the fact that $[\lambda]$ -primitive themes are preserved by $\otimes E_N$ where N is a natural integer, we conclude that $d(E) = d(E^* \otimes E_N)$. Then E admits a $[\lambda]$ -primitive normal sub-theme of rank d . Conversely, if d is the maximal rank of a (normal) $[\lambda]$ -primitive sub-theme of E , then we have $d(E^* \otimes E_N) = d$ and $d(E) = d$. ■

Lemma 4.1.10 *Let E be a rank k $[\lambda]$ -primitive theme and denote by F_j its normal rank j submodule. Let $x \in E \setminus F_{k-1}$. Then the (a,b) -module $\tilde{\mathcal{A}}.x \subset E$ is a rank k theme.*

PROOF. We may assume $E \subset \Xi_\lambda^{(k-1)}$ and then (see [B.10]) we have the equality $F_{k-1} = E \cap \Xi_\lambda^{(k-2)}$. So x contains a non zero term with $(Log s)^{k-1}$ and then the result is clear. ■

Our next lemma shows that the semi-simple filtration of a normal submodule of a fresco E is the trace on this submodule of the semi-simple filtration of E .

Lemma 4.1.11 *Let E be a fresco and F any normal submodule of E . Then for any $j \in \mathbb{N}$ we have $S_j(E) \cap F = S_j(F)$.*

PROOF. By induction on $j \geq 1$. First $S_1(E) \cap F$ is semi-simple in F so contained in $S_1(F)$ by definition. But conversely, $S_1(F)$ is semi-simple, so contained in $S_1(E)$ and also in F .

Let assume now that $j \geq 2$ and that the result is proved for $j-1$. Consider now the quotient $E/S_{j-1}(E)$. As $S_{j-1}(E) \cap F = S_{j-1}(F)$, $F/S_{j-1}(F)$ is a submodule of $E/S_{j-1}(E)$. Now by the case $j=1$ $S_j(F)/S_{j-1}(F)$ which is, by definition, $S_1(F/S_{j-1}(F))$ is equal to $S_1(E/S_{j-1}(E)) \cap (F/S_{j-1}(F))$. So we obtain

$$S_j(F)/S_{j-1}(F) = (S_j(E)/S_{j-1}(E)) \cap (F/S_{j-1}(F)).$$

This implies the equality $S_j(F) = S_j(E) \cap F$. ■

Lemme 4.1.12 *Let $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ be a short exact sequence of frescos. Then we have the inequalities*

$$\sup\{d(F), d(G)\} \leq d(E) \leq d(F) + d(G).$$

PROOF. The inequality $d(F) \leq d(E)$ is obvious from the previous lemma. The inequality $d(G) \leq d(E)$ is then a consequence of the property iv) in proposition 4.1.9.

Now let $\varphi : E \rightarrow \Xi_\lambda$ be an $\tilde{\mathcal{A}}$ -linear map with rank δ . Then the restriction of φ to F has rank $\leq d(F)$. So $\varphi(F)$ is contained in T_d , the normal sub-theme of $\varphi(E)$ of rank $d = d(F)$. The map $\tilde{\varphi} : E/F \rightarrow \Xi_\lambda$ defined by composition of φ with an injection of the theme $\varphi(E)/T_d$ in Ξ_λ has rank $\delta - d \leq d(E/F)$. So the inequality $\delta \leq d(F) + d(E/F)$ is proved. $\blacksquare$

4.2 Co-semi-simple filtration.

Lemme 4.2.1 *Let E be a fresco. Then there exists a normal submodule $\Sigma^1(E)$ which is the minimal normal submodule Σ such that E/Σ is semi-simple.*

PROOF. First recall that if T is a theme and $T \otimes E_\delta$ is geometric for some $\delta \in \mathbb{Q}$, then $T \otimes E_\delta$ is again a theme.

We shall prove that if E is a fresco and if $N \in \mathbb{Z}$ is such that $E \otimes E_N$ is again a fresco, we have the equality of submodules in $E \otimes E_N$:

$$S_1(E \otimes E_N) = S_1(E) \otimes E_N. \quad (@)$$

As $S_1(E) \otimes E_N$ is a normal semi-simple submodule of $E \otimes E_N$ the inclusion $\supset$ in (@) is clear.

Conversely, $S_1(E \otimes E_N) \otimes E_{-N}$ is a semi-simple submodule of $E \simeq E \otimes E_N \otimes E_{-N}$. So we obtain $S_1(E \otimes E_N) \otimes E_{-N} \subset S_1(E)$ and we conclude by tensoring by E_N . Now we shall prove that $S_1(E^* \otimes E_N)^* \otimes E_N$ is a fresco and does not depend of N , large enough.

Let $\lambda_j + j, j \in [1, k]$ the sequence corresponding to the quotient of a J-H. of E . Then let $q \in \mathbb{N}$ such that $\lambda_j + j \in]k, k+q[$ for all $j \in [1, k]$. The corresponding J-H. for $E^* \otimes E_N$ has quotients associated to the numbers $-(\lambda_j + j) + k + N$ which are in $]N - q, N[\subset]k, +\infty[$ for $N > k + q$. So $E^* \otimes E_N$ is a fresco. Then $S_1(E^* \otimes E_N)$ is also a fresco and has a J-H. sequence corresponding to numbers in a subset of the previous ones. Dualizing again, we obtain that $S_1(E^* \otimes E_N)^* \otimes E_N$ has a J-H. sequence with corresponding numbers $-(\mu_j + j) + k + N$ with $\mu_j + j \in]N - q, N[$. So $S_1(E^* \otimes E_N)^* \otimes E_N$ is a fresco which is a quotient of E .

We want to show that this quotient is independant of the choice of N large enough. This is consequence of the fact that

$$\begin{aligned} S_1(E \otimes E_{N+1})^* &= (S_1(E \otimes E_N) \otimes E_1)^* \\ &= S_1(E \otimes E_N)^* \otimes E_{-1} \end{aligned}$$

and so

$$S_1(E \otimes E_{N+1})^* \otimes E_{N+1} = S_1(E \otimes E_N)^* \otimes E_N.$$

As $S_1(E^* \otimes E_N)$ is the maximal semi-simple submodule in $E^* \otimes E_N$, we conclude that $S_1(E^* \otimes E_N)^* \otimes E_N$ is the maximal quotient of E which is semi-simple. So we have $\Sigma^1(E) = \left(E^* \otimes E_N / S_1(E^* \otimes E_N)\right)^* \otimes E_N \subset E$. $\blacksquare$

Définition 4.2.2 Let E be a fresco and define inductively the normal submodules $\Sigma^j(E)$ as follows : $\Sigma^0(E) := E$ and $\Sigma^{j+1}(E) := \Sigma^1(\Sigma^j(E))$. We call $\Sigma^j(E), j \geq 0$ the **co-semi-simple filtration** of E .

Note that Σ^j / Σ^{j+1} is the maximal semi-simple quotient of Σ^j for each j .

Lemme 4.2.3 Let E be a fresco. The normal submodules $\Sigma^j(E)$ satisfies the following properties :

- i) For any $j \in [0, d(E) - 1]$ we have $\Sigma^{j+1}(E) \subset \Sigma^j(E) \cap S_{d-j-1}(E)$ where $d := d(E)$ is the ss-depth of E .
- ii) For any $[\lambda]$ -primitive sub-theme T of rank t in $\Sigma^j(E)$ we have the inclusion $F_{t-p}(T) \subset \Sigma^{j+p}$, where $p \in [0, t]$ and $F_{t-p}(T)$ is the rank $t - p$ normal sub-theme of T .
- iii) Put $d := d(E)$. Then we have $d(\Sigma^j(E)) = d - j$ for each $j \in [0, d]$. This implies that $\Sigma^d(E) = \{0\}$ and that $\Sigma^{d-1}(E) \neq \{0\}$ is semi-simple.
- iv) For any normal submodule $F \subset E$ we have $\Sigma^j(F) \subset \Sigma^j(E) \cap \Sigma^{j-1}(F)$.

REMARKS.

1. The inclusion in i) implies $\Sigma^j(E) \subset S_{d-j}(E) \quad \forall j \in [0, d(E)]$.
2. The filtration $\Sigma^j, j \in [0, d]$, is strictly decreasing because of iii).

PROOF. Let us prove i) by induction on $j \in [0, d(E) - 1]$. As i) is obvious for $j = 0$ assume $j \geq 1$ and i) proved for $j - 1$. So we know that $\Sigma^j(E) \subset S_{d-j}(E)$. The quotient $S_{d-j}(E) / S_{d-j-1}(E)$ is semi-simple, by definition of $S_{d-j}(E)$, and so is its submodule $\Sigma^j(E) / \Sigma^j(E) \cap S_{d-j-1}(E)$. The definition of $\Sigma^{j+1}(E)$ implies then that we have $\Sigma^{j+1}(E) \subset \Sigma^j(E) \cap S_{d-j-1}(E)$. So i) is proved.

To prove ii) it is enough to show it for $p = 1$, by an obvious iteration. By definition $\Sigma^j(E) / \Sigma^{j+1}(E)$ is semi-simple. So is the submodule $T / T \cap \Sigma^{j+1}(E)$. As it is also a $[\lambda]$ -primitive theme, its rank is ≤ 1 showing that $T \cap \Sigma^{j+1}(E)$ contains the corank 1 normal submodule $F_{t-1}(T)$ of T .

To prove iii) let $d := d(E)$ and let T a sub-theme in E of rank d . Then, thanks to ii) with $j = 0$, $\Sigma^1(E) \cap T$ contains a sub-theme of rank $\geq d - 1$. So $d(\Sigma^1(E)) \geq d - 1$. Assume that $d(\Sigma^1(E)) = d$, then we obtain, thanks to

iteration of the previous inequality, that $d(\Sigma^{d-1}(E)) \geq 2$. But from i) we know that $\Sigma^{d-1}(E) \subset S_1(E)$ is semi-simple. This is a contradiction. So we obtain $d(\Sigma^1(E)) = d - 1$ and then $d(\Sigma^j(E)) = d - j$ for each $j \in [0, d]$.

To prove iv) we shall make an induction on $j \in [0, d(E)]$. As the case $j = 0$ is obvious⁸, assume $j \geq 1$ and the case $j - 1$ proved. As $\Sigma^j(E) \cap F / \Sigma^{j+1}(E) \cap F$ is a submodule of $\Sigma^j(E) / \Sigma^{j+1}(E)$ which is semi-simple by definition, it is semi-simple and so we have $\Sigma^1(\Sigma^j(E) \cap F) \subset \Sigma^{j+1}(E) \cap F$. Now to conclude, as we know that $\Sigma^j(F) \subset \Sigma^j(E)$, it is enough to remark that for $G \subset H$ we have $\Sigma^1(G) \subset \Sigma^1(H) \cap G$: as $H / \Sigma^1(H)$ is semi-simple, its submodule $G / G \cap \Sigma^1(H)$ is also semi-simple, and so $\Sigma^1(G)$ is contained in $\Sigma^1(H) \cap G$. ■

REMARK. We shall prove in section 5 that $E / S_1(E)$ and $\Sigma^1(E)$ are rank $d(E) - 1$ themes and that any normal rank $d(E)$ theme in E contains $\Sigma^1(E)$.

4.3 Computation of the ss-depth.

Définition 4.3.1 Let E be a rank k $[\lambda]$ -primitive fresco and consider $[F] := \{F_j, j \in [1, k]\}$ be any J-H. sequence of E . We shall say that $j \in [1, k - 1]$ is a **non commuting index** for $[F]$ if the quotient F_{j+1} / F_{j-1} is a theme. If it is not the case we shall say that j is a **commuting index**. Note that in this case the quotient F_{j+1} / F_{j-1} is semi-simple.

Lemme 4.3.2 Let E be a rank $k \geq 2$ $[\lambda]$ -primitive fresco and let $F_j, j \in [1, k]$ be a J-H. sequence of E . Assume that the $\mathcal{A}$ -linear map $\varphi : F_{k-1} \rightarrow \Xi_\lambda$ has rank $\delta \geq 1$ and that E / F_{k-2} is a theme. Then any $\tilde{\varphi} : E \rightarrow \Xi_\lambda$ extending φ has rank $\delta + 1$.

PROOF. Let e be a generator of E such that $(a - \lambda_{k-1}.b).S_{k-1}^{-1}.(a - \lambda_k.b).e$ is in F_{k-2} . So, if $\lambda_k = \lambda_{k-1} + p_{k-1} - 1$ we have either $p_{k-1} = 0$ or $p_{k-1} \geq 1$ and the coefficient of $b^{p_{k-1}}$ in S_{k-1} does not vanish. Put $\varepsilon := S_{k-1}^{-1}.(a - \lambda_k.b).e$; it is a generator of F_{k-1} . Up to a non zero constant, we may assume that

$$\varphi(\varepsilon) - s^{\lambda_{k-1}-1}.(\text{Log } s)^{\delta-1} / (\delta - 1)! \in \Xi_\lambda^{(\delta-2)}.$$

Now we want to define $\tilde{\varphi}(e) = x$ where x is a solution in Ξ_λ of the equation

$$(a - \lambda_k.b).x = S_{k-1}.\varphi(e).$$

Then it is easy to find that, because the coefficient of $b^{p_{k-1}}$ in S_{k-1} is not zero, we have

$$x - s^{\lambda_{k-1}-1}.(\text{Log } s)^\delta / \delta! \in \Xi_\lambda^{(\delta-1)}.$$

Now the degree in $\text{Log } s$ gives our assertion. ■

⁸with the convention $\Sigma^{-1}(G) := G$.

Corollaire 4.3.3 *In the situation of the lemma 4.3.2 we have the equality $d(E) = d(F_{k-1}) + 1$.*

Lemme 4.3.4 *Let E be a rank k $[\lambda]$ -primitive fresco and $[F] := \{F_j, j \in [1, k]\}$, be any J-H. sequence of E . Let $nci(F)$ be the number of **non commuting indices** for the J-H. sequence $[F]$. Then we have the inequality*

$$d(E) \geq nci(F) + 1$$

PROOF. We shall prove this by induction on the rank of E . The cases of rank 1 and 2 are clear. Let assume $k \geq 3$ and the inequality proved in rank $\leq k - 1$. Consider a J-H. sequence $F_j, j \in [1, k]$, for E and assume first that E/F_{k-2} is semi-simple. Then we have, denoting $[G]$ the J-H. sequence $\{F_j, j \in [1, k - 1]\}$, for F_{k-1} :

$$nci(F) = nci(G) \leq d(F_{k-1}) - 1 \leq d(E) - 1$$

using the induction hypothesis and lemma 4.1.12 ; it concludes this case.

Assume now that E/F_{k-2} is a theme. Then using corollary 4.3.3 we have $d(E) = d(F_{k-1}) + 1$. So we get using again the inductive hypothesis :

$$nci(F) = nci(G) + 1 \leq d(F_{k-1}) = d(E) - 1$$

which concludes the proof. ■

REMARKS.

1. This inequality may be strict for several J-H. sequences, including the principal one : there are examples of rank 3 fresco with a principal J-H. sequence $[F]$ such that $nci(F) = 0$ which are not semi-simple (see 3.3).
2. We shall see using the corollary of the theorem 5.1.1 (see the remark following 5.1.2) that for any fresco E there always exists a J-H. sequence $[F]$ for which we have the equality $d(E) = nci(F) + 1$.

4.4 Embedding for a semi-simple fresco.

The aim of this paragraph is to prove the following embedding theorem for semi-simple $[\lambda]$ -primitive frescos.

Proposition 4.4.1 *Let E be a rank k semi-simple $[\lambda]$ -primitive fresco. Then there exists an $\tilde{A}$ -linear injective map $\varphi : E \rightarrow \Xi_\lambda \otimes \mathbb{C}^l$ if and only if $l \geq k$.*

PROOF. To show that the existence of $\varphi : E \rightarrow \Xi_\lambda \otimes \mathbb{C}^l$ implies $l \geq k$, remark that for any linear form $\alpha : \mathbb{C}^l \rightarrow \mathbb{C}$ the composed map $(1 \otimes \alpha) \circ \varphi$ has rank at most 1. So the inequality $l \geq k$ is clear. To prove that there exists an $\tilde{\mathcal{A}}$ -linear injective map from E to $\Xi_\lambda \otimes \mathbb{C}^k$ we shall use the following lemma.

Lemme 4.4.2 *Let $\lambda_1, \dots, \lambda_k, k \geq 2$, be numbers in $[\lambda] \in \mathbb{Q}/\mathbb{Z}$ such that $\lambda_{j+1} = \lambda_j + p_j - 1$ for each $j \in [1, k-1]$ with $p_j < 0$. Put*

$$Q := (a - \lambda_2.b).S_2^{-1} \dots S_{k-1}^{-1}.(a - \lambda_k.b) \quad \text{and} \quad P := (a - \lambda_1.b).S_1^{-1}.Q$$

where $S_j, j \in [1, k-1]$ are invertible elements in $\mathbb{C}[[b]]$. Assume also that $\lambda_1 > k-1$. Then there exists a unique element $T \in \mathbb{C}[[b]]$ which satisfies

$$Q.T.s^{\lambda_1-k} = S_1.s^{\lambda_1-1}.$$

Moreover T is invertible in $\mathbb{C}[[b]]$.

PROOF. We begin by the proof of the case $k = 2$. Then we look for $T \in \mathbb{C}[[b]]$ such that $(a - \lambda_2.b).T.s^{\lambda_1-2} = S_1.s^{\lambda_1-1}$. This equation is equivalent to the differential equation

$$b.T' - p_1.T = (\lambda_1 - 1).S_1$$

which has an unique solution in $\mathbb{C}[[b]]$ for any $S_1 \in \mathbb{C}[[b]]$ because $p_1 < 0$. Moreover, we have $-p_1.T(0) = (\lambda_1 - 1).S_1(0)$, so T is invertible as S_1 is invertible.

Let now prove the lemma by induction on $k \geq 2$. We may assume $k \geq 3$ and the lemma proved for $k-1$. Put $Q = (a - \lambda_2.b).S_2^{-1}.R$. Our equation is

$$(a - \lambda_2.b).S_2^{-1}.R.T.s^{\lambda_1-k} = S_1.s^{\lambda_1-1}. \quad (@)$$

Remark that $S_2^{-1}.R.T.s^{\lambda_1-k} = V.s^{\lambda_1-2}$ for some $V \in \mathbb{C}[[b]]$. So, let $U \in \mathbb{C}[[b]]$ the unique solution of the equation

$$(a - \lambda_2.b).U.s^{\lambda_1-2} = S_1.s^{\lambda_1-1}$$

and consider now the equation in $T \in \mathbb{C}[[b]]$:

$$R.T.s^{\lambda_1-k} = S_2.U.s^{\lambda_1-2}. \quad (@@)$$

The inductive hypothesis shows that there exists a unique invertible $T \in \mathbb{C}[[b]]$ which is solution of (@@). Then it satisfies (@). The uniqueness of the solution T of (@) is consequence of the uniqueness of U and uniqueness in the inductive hypothesis. ■

END OF THE PROOF OF PROPOSITION 4.4.1. We shall prove the assertion by induction on k the rank of E . The case $k = 1$ is obvious, so we may assume $k \geq 2$ and that we have a $\tilde{\mathcal{A}}$ -linear injective map $\varphi : F \rightarrow \Xi_\lambda \otimes \mathbb{C}^{k-1}$ for a semi-simple $[\lambda]$ -primitive fresco F of rank $k-1$. Now using the notations of the previous lemma and proposition 4.1.4 we may assume that $E := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.P$ and put $F := \tilde{\mathcal{A}}/\tilde{\mathcal{A}}.Q$. Let e be a generator of E with annihilator $\tilde{\mathcal{A}}.P$. It induces a generator $[e]$ of F with annihilator $\tilde{\mathcal{A}}.Q$. Note that $F = E/E_{\lambda_1}$ where $E_{\lambda_1} \subset E$ is generated by $Q.e$. So $\varphi([e]) \in \Xi_\lambda \otimes \mathbb{C}^{k-1}$ is killed by Q . Using the previous lemma, we find $T \in \mathbb{C}[[b]]$ such that $Q.T.s^{\lambda_1-k} = S_1.s^{\lambda_1-1}$. So $T.s^{\lambda_1-k}$ is killed by $P = (a - \lambda_1.b).S_1^{-1}.Q$. Define now $\psi : E \rightarrow \Xi_\lambda$ via $\psi(e) := T.s^{\lambda_1-k}$ and also $\Phi : E \rightarrow \Xi_\lambda \otimes \mathbb{C}^k$ as the direct sum of $\varphi \circ q$ and ψ , where $q : E \rightarrow F$ is the obvious quotient map given by $e \mapsto [e]$. Then Φ is injective : if $\Phi(x) = 0$ then $q(x) = 0$ so x is in $\text{Ker } q := \mathbb{C}[[b]].Q(e) \simeq E_{\lambda_1}$. Then

$$\psi(x) = \psi(X.Q.e) = X.Q.T.s^{\lambda_1-k} = X.S_1.(S_1^{-1}.Q.T).s^{\lambda_1-k} = X.S_1.s^{\lambda_1-1} = 0$$

if $x := X.Q(e)$ with $X \in \mathbb{C}[[b]]$. As $Q.e$ generates a free (rank 1) $\mathbb{C}[[b]]$ -module and S_1 is invertible in $\mathbb{C}[[b]]$, we conclude that $X = 0$ and also $x = 0$. ■

5 Quotient themes of a $[\lambda]$ -primitive fresco.

5.1 Structure theorem for $[\lambda]$ -primitive frescos.

Now we are ready to describe the precise structure of a $[\lambda]$ -primitive fresco. We shall then deduce the possible ($[\lambda]$ -primitive) quotient themes of a any given $[\lambda]$ -primitive fresco.

Théorème 5.1.1 *Let E be a $[\lambda]$ -primitive fresco and let $d := d(E)$ be its ss-depth. Then there exists a J-H. sequence $G_j, j \in [1, k]$ for E with the following properties :*

- i) *The quotient E/G_{k-d} is a theme (with rank d).*
- ii) *We have the equality $G_{k-d+1} = S_1(E)$.*

REMARK. By twisted duality we obtain also a J-H. sequence $G'_j, j \in [1, k]$, such that $G'_{d-1} = \Sigma^1(E)$ and G'_d is a theme ; then E/G'_{d-1} is the maximal semi-simple quotient of E .

PROOF. Let $\varphi : E \rightarrow \Xi_\lambda$ an $\tilde{\mathcal{A}}$ -linear map with rank d . Denote $F_j(\varphi)$ the J-H. sequence of the $[\lambda]$ -primitive theme $\varphi(E)$. Put $H_j := \varphi^{-1}(F_j(\varphi))$ for $j \in [0, d]$. Note that $H_0 = \text{Ker } \varphi$. We shall show that $H_1 = S_1(E)$.

To show that H_1 is semi-simple, assume that we have a rank 2 $\tilde{\mathcal{A}}$ -linear map $\theta : H_1 \rightarrow \Xi_\lambda$. Then $E/\text{Ker } \theta$ has the following J-H. sequence

$$0 \subset \theta^{-1}(F_1(\theta))/\text{Ker } \theta \subset H_1/\text{Ker } \theta \subset \cdots \subset H_d/\text{Ker } \theta.$$

As $H_1/\text{Ker } \theta$ and $(H_{j+1}/\text{Ker } \theta)/(H_{j-1}/\text{Ker } \theta) \simeq H_{j+1}/H_{j-1}$ are $[\lambda]$ -primitive rank 2 themes for j in $[1, d-1]$, this would imply that $E/\text{Ker } \theta$ is a rank $d+1$ $[\lambda]$ -primitive theme, contradicting the definition of d .

Then H_1 is semi-simple. But as $S_1(E)$ is contained in $H_1 := \varphi^{-1}(F_1(\varphi))$ the equality $H_1 = S_1(E)$ is proved.

Define now $G_{k-d+j} := H_j$ for $j \in [0, d]$, and complete the J-H. sequence

$G_j, j \in [1, k]$ of E by choosing a J-H. sequence $G_j, j \in [1, k-d]$, for H_0 . ■

We have the following easy consequences of this theorem.

Corollaire 5.1.2 *Let E be a $[\lambda]$ -primitive fresco and let $d := d(E)$ be its ss-depth. Then $E/S_1(E)$ and $\Sigma^1(E)$ are $[\lambda]$ -primitive themes of rank $d-1$ and we have*

$$\text{rk}(S_1(E)) + d(E) = \text{rk}(E) + 1. \quad \text{and} \quad \text{rk}(\Sigma^1(E)) = d(E) - 1.$$

For each $j \in [2, d]$ the rank of $S_j(E)/S_{j-1}(E)$ is 1. Moreover, any rank d quotient theme T of E satisfies $E/S_1(E) \simeq T/F_1(T)$. Dually, any rank d theme contains $\Sigma^1(E)$.

PROOF. With the notations of the theorem, let $T := E/G_{k-d}$. Then G_{k-d+1}/G_{k-d} is $F_1(T)$ the unique rank 1 normal submodule of the $[\lambda]$ -primitive theme T . Then we obtain that $E/S_1(E) \simeq E/G_{k-d+1} \simeq T/F_1(T)$ proving our first assertion. The computation of the rank of $S_1(E)$ follows.

Consider now the exact sequence

$$0 \rightarrow S_1(E) \rightarrow E \rightarrow T(E) \rightarrow 0$$

where $T(E)$ is the $d-1$ $[\lambda]$ -primitive theme $E/S_1(E)$. Dualizing and tensoring by E_N for N a large enough integer gives the exact sequence

$$0 \rightarrow T(E)^* \otimes E_N \rightarrow E^* \otimes E_N \rightarrow S_1(E)^* \otimes E_N \rightarrow 0$$

where $S_1(E)^* \otimes E_N$ is semi-simple. This implies that $\Sigma^1(E^* \otimes E_N) \subset T(E)^* \otimes E_N$. And we have equality because we know that $E^* \otimes E_N$ contains a rank d theme, so the dimension of $\Sigma^1(E^* \otimes E_N)$ is at most $d(E) - 1$. Then we conclude that $\Sigma^1(E)$ is a theme and that its rank is $d(E) - 1$.

We know that $E/S_1(E)$ is a theme, so $S_1(E/S_1(E)) = F_1(E/S_1(E))$ is rank 1 for $d(E) \geq 2$. A similar argument shows that $S_j(E)/S_{j-1}(E)$ for each $j \in [2, d]$. In fact it is naturally isomorphic to $F_{j-1}(T)/F_{j-2}(T)$ where $T := T(E)$.

Let T' any rank d quotient theme of E . As E is $[\lambda]$ -primitive, so is T' and we may assume that $T' := \varphi(E)$ where $\varphi \in \text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda)$. But now $S_1(E)$ is in $\varphi^{-1}(F_1(T'))$, so we have a surjective map, induced by φ :

$$\tilde{\varphi} : E/S_1(E) \rightarrow T'/F_1(T')$$

between two $[\lambda]$ -primitive themes of the same rank $d - 1$. This must be an isomorphism. ■

REMARK. Building a J-H. sequence of E via the exact sequence

$$0 \rightarrow S_1(E) \rightarrow E \rightarrow T(E) \rightarrow 0$$

we find that the number of non commuting indices in such a J-H. sequence is exactly $d(E) - 1$. Of course we may put the non commuting indices at the beginning by using the exact sequence

$$0 \Sigma^1(E) \rightarrow E \rightarrow E/\Sigma^1(E) \rightarrow 0.$$

5.2 Embedding dimension for a $[\lambda]$ -primitive fresco.

From the embedding result in the semi-simple case 4.4.1 and the structure theorem 5.1.1 we shall deduce precise embedding theorem for $[\lambda]$ -primitive frescos.

Proposition 5.2.1 *Let E be a $[\lambda]$ -primitive fresco. Then there exists an injective $\tilde{\mathcal{A}}$ -linear map $\varphi : E \rightarrow \Xi_\lambda \otimes \mathbb{C}^l$ if and only if $l \geq rk(E) - d(E) + 1$.*

PROOF. If we have such a φ its restriction to $S_1(E)$ is an embedding and so $l \geq rk(S_1(E)) = rk(E) - d(E) + 1$.

Conversely, we shall prove that there exists an embedding of E in $\Xi_\lambda \otimes \mathbb{C}^l$ with $l = rk(E) - d(E) + 1$. By the proposition 4.4.1 we may begin with an injective $\tilde{\mathcal{A}}$ -linear map $\varphi : S_1(E) \rightarrow \Xi_\lambda \otimes \mathbb{C}^l$ with $l := rk(E) - d(E) + 1$. Put $\varphi := \oplus_{i=1}^l \varphi_i$ where $\varphi_i : S_1(E) \rightarrow \Xi_\lambda$. Now, for each $i \in [1, l]$ we may find an extension $\Phi_i : E \rightarrow \Xi_\lambda$ to φ_i thanks to the surjectivity of $\text{Hom}_{\tilde{\mathcal{A}}}(E, \Xi_\lambda) \rightarrow \text{Hom}_{\tilde{\mathcal{A}}}(S_1(E), \Xi_\lambda)$ (see section 2.1). Then we may define $\Phi := \oplus_{i \in [1, l]} \Phi_i : E \rightarrow \Xi_\lambda \otimes \mathbb{C}^l$ which is an extension of φ to E . Moreover, such an extension is injective because its kernel cannot meet non trivially $S_1(E)$ and so does not contain any normal rank 1 submodule of E . So the kernel has to be $\{0\}$. ■

5.3 Quotient themes of a $[\lambda]$ -primitive fresco.

Now we shall describe all quotient themes of a given $[\lambda]$ -primitive fresco. We begin by the description of quotient themes of maximal rank.

Proposition 5.3.1 *Let E be a $[\lambda]$ -primitive fresco of rank k . Then any rank $d := d(E)$ quotient theme of E is obtained as follows : let K be a corank 1 normal submodule of $S_1(E)$ and assume that $K \cap L(E) = \{0\}$ where $L(E) := \Sigma^1(S_2(E))$. Then E/K is a rank d theme.*

PROOF. Consider $K \subset S_1(E)$ a corank 1 normal submodule in $S_1(E)$ such that $K \cap L(E) = \{0\}$. By definition of $L(E)$ the quotient $S_2(E)/K$ is rank 2 and not semi-simple. So it is a theme. We have the following Jordan-Hölder sequence for E/K :

$$0 \subset S_1(E)/K \subset S_2(E)/K \subset \cdots \subset S_d(E)/K = E/K \quad (@)$$

But $S_2(E)/K$ and each $S_{j+2}(E)/S_j(E)$ for $j \in [1, d-2]$ is a theme of rank 2. So from [B.10] we conclude that E/K is a rank d theme.

Conversely, if E/K is a rank d theme, consider $S_1(E)/S_1(E) \cap K \hookrightarrow E/K$. As $S_1(E)/S_1(E) \cap K$ is semi-simple and E/K is a theme, the rank of $S_1(E)/S_1(E) \cap K$ is at most 1. It is not 0 because K has rank $k-d$ and $S_1(E)$ has rank $k-d+1$. So K is contained in $S_1(E)$ and has corank 1 in it. If K contains $L(E)$ then $S_2(E)/K$ is semi-simple and of rank 2, if we assume $d \geq 2$. But it is contained in E/K which is a theme, so we get a contradiction.

For $d = 1$ in the previous proposition (so E semi-simple) we have $S_2(E) = E$ so $L(E) = \{0\}$ and any corank 1 normal submodule of $S_1(E) = E$ gives a rank 1 quotient which is, of course a rank 1 theme. ■

In the statement of the theorem we shall denote by L_j for $j \in [1, d-1]$ the rank j theme defined as $\Sigma^1(S_{j+1}(E))$. So, by definition, $S_{j+1}(E)/\Sigma_j$ is semi-simple, and L_j is a normal submodule which is minimal for this property. We have seen that L_j is then a theme with rank $d(S_{j+1}(E)) - 1$; as we know that $d(S_{j+1}(E)) = j+1$, the rank of L_j is j . In fact we have $L_j = F_j(\Sigma^1(E))$ for each $j \in [0, d-1]$ and so $L_1 = L(E)$.

Théorème 5.3.2 *Let E be a $[\lambda]$ -primitive fresco of rank k . Put $d := d(E)$ and denote by $S_j, j \in [1, d]$ the semi-simple filtration of E . Assume that $d \geq 2$ and let Σ_j be the first term of the co-semi-simple filtration of S_{j+1} .*

Let $K \subset E$ be a normal submodule such that E/K is a theme. Then we have the following possibilities :

1. *If K contains $S_1(E)$, then E/K is a quotient theme of the rank $d-1$ theme $E/S_1(E)$ and we have exactly one such quotient for each rank in $[1, d-1]$.*
2. *If $K \cap L_1 = \{0\}$, then E/K is a quotient of the rank d theme $E/K \cap S_1(E)$ which belongs to the quotient themes described in the previous proposition.*
3. *If K contains L_{j_0} but not L_{j_0+1} (so $K \cap \Sigma^1(E)$ has rank j_0), we may apply the previous case to $E' := E/L_{j_0}$ and $K' := K/L_{j_0}$ and find that $E/K = E'/K'$ is a quotient of the rank $d-j_0$ quotient theme $E'/K' \cap S_1(E')$.*

In this situation we have $S_1(E') = S_{j_0+1}(E)/L_{j_0}$ and $d(E') = d - j_0$, with $rk(E') = rk(E) - j_0$. So the rank of E/K is at most $d - j_0$.

REMARKS.

- i) The case 2 of the previous theorem is the case 3 with $j_0 = 0$. We emphasize on this case because the rank 1 quotients theme is the most interesting case.
- ii) Let $\varphi : E \rightarrow T$ be a surjective $\tilde{\mathcal{A}}$ -linear map on a rank $\delta \geq 2$ theme T . Then $S_1(E) \subset \varphi^{-1}(F_1(T))$ and so the map φ induces a surjection $E/S_1(E) \rightarrow T/F_1(T)$. As $E/S_1(E)$ is a $[\lambda]$ -primitive theme, it has an unique quotient of rank $\delta - 1$. So the quotient theme $T/F_1(T)$ depends only on δ and E , not on T . In the case $\delta = d(E)$ we find that $T/F_1(T) \simeq E/S_1(E)$ for any choice of T .
- iii) The previous theorem gives very few information on the rank 1 quotients, because any corank 1 normal submodule contains $\Sigma^!(E)$. They will be described in the next proposition.

PROOF. The first case is clear.

Assume that $L_1 \cap K = \{0\}$; then K does not contain S_1 . But $S_1/K \cap S_1$ is semi-simple and is contained in the theme E/K . So it has rank ≤ 1 . As we know that S_1 is not contained in K , the rank is exactly 1, and $K \cap S_1$ has corank 1 in S_1 . As $K \cap L_1 = \{0\}$ the previous proposition shows that $E/K \cap S_1$ is a rank d theme. So the case 2 is proved.

For the proof of the case 3 it is enough to prove the equalities

$$\begin{aligned} S_1(E') &= S_{j_0+1}(E)/L_{j_0} & L_1(E') &= L_{j_0+1}/L_{j_0} & \text{and} \\ d(E') &= d - j_0, & rk(E') &= rk(E) - j_0. \end{aligned}$$

As we know that L_j is a theme of rank j , because $d(S_{j+1}) = j + 1$, the rank of E' is $rk(E) - j_0$. The equality $d(L_{j_0}) = j_0$ is then clear.

As $S_{j_0+1}(E)/L_{j_0}$ is semi-simple, we have $S_{j_0+1}(E)/L_{j_0} \subset S_1(E')$. But now we know that they have same rank because

$$rk(S_1(E')) = rk(E') - d(E') + 1 = rk(E) - j_0 - (d - j_0) + 1 = rk(E) - d + 1.$$

Also $L_1(E')$ and L_{j_0+1}/L_{j_0} are rank 1 and normal submodules of E' . And as $S_2(E') = S_{j_0+2}/L_{j_0}$, we have $S_2(E')/[L_{j_0+1}/L_{j_0}] = S_{j_0+2}/S_{j_0+1}$ is semi-simple (rank 1), this gives the inclusion

$$L_1(E') := \Sigma^1(S_2(E')) \subset L_{j_0+1}/L_{j_0}$$

and so the equality $L_1(E') = L_{j_0+1}/L_{j_0}$ is proved. ■

Proposition 5.3.3 (The rank 1 quotients) *Let E be a $[\lambda]$ -primitive rank k fresco. Put $d := d(E)$. Then any rank 1 quotient of E is a rank 1 quotient of $E/\Sigma^1(E)$. As $E/\Sigma^1(E)$ is semi-simple of rank $k - d + 1$ it shows that there are exactly $k - d + 1$ isomorphism classes of such a rank 1 quotient and they corresponds to the fundamental invariants of $E/\Sigma^1(E)$ as follows : if $\lambda_1, \dots, \lambda_{k-d+1}$ are numbers associated to any J-H. sequence of $E/\Sigma^1(E)$, the isomorphism classes of rank 1 quotients of E are given by*

$$\lambda_1 - k + d, \dots, \lambda_2 - k + d - 1, \dots, \lambda_{k-d+1}.$$

PROOF. Let H be a normal co-rank 1 submodule of E . As E/H is semi-simple, H contains $\Sigma^1(E)$ so E/H is a rank 1 quotient of $E/\Sigma^1(E)$. The converse is obvious. ■

REMARK. Assume $d(E) \geq 2$. With the exception of the (unique) rank 1 quotient of $E/S_1(E)$ which is $E/S_{d-1}(E)$, no rank 1 quotient of E may be the rank 1 quotient of a quotient theme of rank ≥ 2 of E . Another way to say that is the following : for any rank $r \geq 2$ quotient theme T of E we have $T/F_{r-1}(T) \simeq E/S_{d-1}(E)$.

EXEMPLE. Let E a rank 3 $[\lambda]$ -primitive fresco with $d(E) = 2$ (so E is not semi-simple and is not a theme). Then there exists $k - d + 1 = 2$ isomorphism classes of rank 1 quotients of E .

For instance assume that we have a J-H. sequence $0 \subset F_1 \subset F_2 \subset F_3 = E$, with $F_2 = S_1(E)$ and such E/F_1 is a rank 2 theme.

If $E \simeq (a - \lambda_1.b)S_1^{-1}.(a - \lambda_2.b)S_2^{-1}.(a - \lambda_3.b)$ this means, with $\lambda_{j+1} = \lambda_j + p_j - 1$ for $j = 1, 2$, that $p_1 < 0$ or $p_1 \geq 1$ and no term in b^{p_1} in S_1 and $p_2 \geq 0$ with a non zero term in b^{p_2} in S_2 .

So we put $F_j/F_{j-1} \simeq E_{\lambda_j}$ for $j \in [1, 3]$; then the rank 1 quotients are isomorphic to $E_{\lambda_3} \simeq E/S_1(E)$ or E_{λ_1-2} , because using the computations of section 3.3 we see that $\Sigma^1(E) \simeq E_{\lambda_2+1}$, and so E_{λ_1-2} is a rank 1 quotient of $E/\Sigma^1(E)$ and a fortiori of E .

To conclude we give a method to compute $L(E)$ in many cases.

Lemme 5.3.4 *Let E be a $[\lambda]$ -primitive fresco and assume that $E \simeq \tilde{A}/\tilde{A}.P$ where*

$$P := (a - \lambda_1.b).S_1^{-1} \dots S_{k-1}^{-1}.(a - \lambda_k.b)$$

where $\lambda_j + j$ is an increasing sequence. Assume that the first non commuting index is $h \in [1, k - 1]$. Then E has a normal sub-theme of rank 2 with fundamental invariants $\lambda_1, \lambda_{h+1} + h$.

PROOF. It is a simple application of the corollary 3.2.2 which also allow to compute the parameter of the rank 2 obtain by commuting in P from the parameter of the rank 2 theme $Fh + 1/F_{h-1}$ and the integers $p_1, \dots, p_h$. ■

REMARKS.

- i) The hypothesis of the lemma means that $p_1, \dots, p_{h-1}$ are positive and that for each $j \in [1, h-1]$ the coefficient of b^{p_j} in S_j is zero. But the coefficient of b^{p_h} in S_h is not zero (and $p_h = 0$ is allowed).
- ii) The lemma implies that $L(E) \simeq E_{\lambda_1}$ is the first term of the principal J-H. sequence of E .

6 Appendix : the existence theorem.

The aim of this appendix is to prove the the following existence theorem for the fresco associated to a relative de Rham cohomology class :

Théorème 6.0.5 *Let X be a connected complex manifold of dimension $n+1$ where n is a natural integer, and let $f : X \rightarrow D$ be an non constant proper holomorphic function on an open disc D in $\mathbb{C}$ with center 0. Let us assume that df is nowhere vanishing outside of $X_0 := f^{-1}(0)$.*

Let ω be a $\mathcal{C}^\infty - (p+1)$ -differential form on X such that $d\omega = 0 = df \wedge \omega$. Denote by E the geometric $\tilde{\mathcal{A}}$ -module $\mathbb{H}^{p+1}(X, (\hat{K}^\bullet, d^\bullet))$ and $[\omega]$ the image of ω in $E/B(E)$. Then $\tilde{\mathcal{A}}.[\omega] \subset E/B(E)$ is a fresco.

Note that this result is an obvious consequence of the finiteness theorem 6.3.4 that we shall prove below. It gives the fact that E is naturally an $\tilde{\mathcal{A}}$ -module which is of finite type over the subalgebra $\mathbb{C}[[b]]$ of $\tilde{\mathcal{A}}$, and so its b -torsion $B(E)$ is a finite dimensional $\mathbb{C}$ -vector space. Moreover, the finiteness theorem asserts that $E/B(E)$ is a geometric (a,b)-module.

6.1 Preliminaries.

Here we shall complete and precise the results of the section 2 of [B.II]. The situation we shall consider is the following : let X be a connected complex manifold of dimension $n+1$ and $f : X \rightarrow \mathbb{C}$ a non constant holomorphic function such that $\{x \in X / df = 0\} \subset f^{-1}(0)$. We introduce the following complexes of sheaves supported by $X_0 := f^{-1}(0)$

1. The formal completion "in f " $(\hat{\Omega}^\bullet, d^\bullet)$ of the usual holomorphic de Rham complex of X .
2. The sub-complexes $(\hat{K}^\bullet, d^\bullet)$ and $(\hat{I}^\bullet, d^\bullet)$ of $(\hat{\Omega}^\bullet, d^\bullet)$ where the subsheaves $\hat{K}^p$ and $\hat{I}^{p+1}$ are defined for each $p \in \mathbb{N}$ respectively as the kernel and the image of the map

$$\wedge df : \hat{\Omega}^p \rightarrow \hat{\Omega}^{p+1}$$

given par exterior multiplication by df . We have the exact sequence

$$0 \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow (\hat{\Omega}^\bullet, d^\bullet) \rightarrow (\hat{I}^\bullet, d^\bullet)[+1] \rightarrow 0. \quad (1)$$

Note that $\hat{K}^0$ and $\hat{I}^0$ are zero by definition.

3. The natural inclusions $\hat{I}^p \subset \hat{K}^p$ for all $p \geq 0$ are compatible with the differential d . This leads to an exact sequence of complexes

$$0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0. \quad (2)$$

4. We have a natural inclusion $f^*(\hat{\Omega}_{\mathbb{C}}^1) \subset \hat{K}^1 \cap \text{Ker } d$, and this gives a sub-complex (with zero differential) of $(\hat{K}^\bullet, d^\bullet)$. As in [B.07], we shall consider also the complex $(\tilde{K}^\bullet, d^\bullet)$ quotient. So we have the exact sequence

$$0 \rightarrow f^*(\hat{\Omega}_{\mathbb{C}}^1) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow (\tilde{K}^\bullet, d^\bullet) \rightarrow 0. \quad (3)$$

We do not make the assumption here that $f = 0$ is a reduced equation of X_0 , and we do not assume that $n \geq 2$, so the cohomology sheaf in degree 1 of the complex $(\hat{K}^\bullet, d^\bullet)$, which is equal to $\hat{K}^1 \cap \text{Ker } d$ does not coincide, in general with $f^*(\hat{\Omega}_{\mathbb{C}}^1)$. So the complex $(\tilde{K}^\bullet, d^\bullet)$ may have a non zero cohomology sheaf in degree 1.

Recall now that we have on the cohomology sheaves of the following complexes $(\hat{K}^\bullet, d^\bullet), (\hat{I}^\bullet, d^\bullet), ([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ and $f^*(\hat{\Omega}_{\mathbb{C}}^1), (\tilde{K}^\bullet, d^\bullet)$ natural operations a and b with the relation $a.b - b.a = b^2$. They are defined in a naïve way by

$$a := \times f \quad \text{and} \quad b := \wedge df \circ d^{-1}.$$

The definition of a makes sens obviously. Let me precise the definition of b first in the case of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ with $p \geq 2$: if $x \in \hat{K}^p \cap \text{Ker } d$ write $x = d\xi$ with $\xi \in \hat{\Omega}^{p-1}$ and let $b[x] := [df \wedge \xi]$. The reader will check easily that this makes sens. For $p = 1$ we shall choose $\xi \in \hat{\Omega}^0$ with the extra condition that $\xi = 0$ on the smooth part of X_0 (set theoretically). This is possible because the condition $df \wedge d\xi = 0$ allows such a choice : near a smooth point of X_0 we can choose coordinnates such $f = x_0^k$ and the condition on ξ means independance of $x_1, \dots, x_n$. Then ξ has to be (set theoretically) locally constant on X_0 which is locally connected. So we may kill the value of such a ξ along X_0 .

The case of the complex $(\hat{I}^\bullet, d^\bullet)$ will be reduced to the previous one using the next lemma.

Lemme 6.1.1 *For each $p \geq 0$ there is a natural injective map*

$$\tilde{b} : \mathcal{H}^p(\hat{K}^\bullet, d^\bullet) \rightarrow \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$$

which satisfies the relation $a.\tilde{b} = \tilde{b}.(b + a)$. For $p \neq 1$ this map is bijective.

PROOF. Let $x \in \hat{K}^p \cap \text{Ker } d$ and write $x = d\xi$ where $x \in \hat{\Omega}^{p-1}$ (with $\xi = 0$ on X_0 if $p = 1$), and set $\tilde{b}([x]) := [df \wedge \xi] \in \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. This is independant on the choice of ξ because, for $p \geq 2$, adding $d\eta$ to ξ does not modify the result as $[df \wedge d\eta] = 0$. For $p = 1$ remark that our choice of ξ is unique.

This is also independant of the the choice of x in $[x] \in \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ because adding $\theta \in \hat{K}^{p-1}$ to ξ does not change $[df \wedge \xi]$.

Assume $\tilde{b}([x]) = 0$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$; this means that we may find $\alpha \in \hat{\Omega}^{p-2}$ such $df \wedge \xi = df \wedge d\alpha$. But then, $\xi - d\alpha$ lies in $\hat{K}^{p-1}$ and $x = d(\xi - d\alpha)$ shows that $[x] = 0$. So $\tilde{b}$ is injective.

Assume now $p \geq 2$. If $df \wedge \eta$ is in $\hat{I}^p \cap \text{Ker } d$, then $df \wedge d\eta = 0$ and $y := d\eta$ lies in $\hat{K}^p \cap \text{Ker } d$ and defines a class $[y] \in \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ whose image by $\tilde{b}$ is $[df \wedge \eta]$. This shows the surjectivity of $\tilde{b}$ for $p \geq 2$.

For $p = 1$ the map $\tilde{b}$ is not surjective (see the remark below).

To finish the proof let us to compute $\tilde{b}(a[x] + b[x])$. Writing again $x = d\xi$, we get

$$a[x] + b[x] = [f.d\xi + df \wedge \xi] = [d(f.\xi)]$$

and so

$$\tilde{b}(a[x] + b[x]) = [df \wedge f.\xi] = a.\tilde{b}([x])$$

which concludes the proof. ■

Denote by $i : (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet)$ the natural inclusion and define the action of b on $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ by $b := \tilde{b} \circ \mathcal{H}^p(i)$. As i is a -linear, we deduce the relation $a.b - b.a = b^2$ on $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ from the relation of the previous lemma.

The action of a on the complex $([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ is obvious and the action of b is zero.

The action of a and b on $f^*(\hat{\Omega}_{\mathbb{C}}^1) \simeq E_1 \otimes \mathbb{C}_{X_0}$ are the obvious one, where E_1 is the rank 1 (a,b)-module with generator e_1 satisfying $a.e_1 = b.e_1$ (or, equivalently, $E_1 := \mathbb{C}[[z]]$ with $a := \times z$, $b := \int_0^z$ and $e_1 := 1$).

Remark that the natural inclusion $f^*(\hat{\Omega}_{\mathbb{C}}^1) \hookrightarrow (\hat{K}^\bullet, d^\bullet)$ is compatible with the actions of a and b . The actions of a and b on $\mathcal{H}^1(\hat{K}^\bullet, d^\bullet)$ are simply induced by the corresponding actions on $\mathcal{H}^1(\hat{K}^\bullet, d^\bullet)$.

REMARK. The exact sequence of complexes (1) induces for any $p \geq 2$ a bijection

$$\partial^p : \mathcal{H}^p(\hat{I}^\bullet, d^\bullet) \rightarrow \mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$$

and a short exact sequence

$$0 \rightarrow \mathbb{C}_{X_0} \rightarrow \mathcal{H}^1(\hat{I}^\bullet, d^\bullet) \xrightarrow{\partial^1} \mathcal{H}^1(\hat{K}^\bullet, d^\bullet) \rightarrow 0 \quad (@)$$

because of the de Rham lemma. Let us check that for $p \geq 2$ we have $\partial^p = (\tilde{b})^{-1}$ and that for $p = 1$ we have $\partial^1 \circ \tilde{b} = Id$. If $x = d\xi \in \hat{K}^p \cap \text{Ker } d$ then $\tilde{b}([x]) = [df \wedge \xi]$ and $\partial^p[df \wedge \xi] = [d\xi]$. So $\partial^p \circ \tilde{b} = Id \quad \forall p \geq 0$. For $p \geq 2$ and

$df \wedge \alpha \in \hat{I}^p \cap \text{Ker } d$ we have $\partial^p[df \wedge \alpha] = [d\alpha]$ and $\tilde{b}[d\alpha] = [df \wedge \alpha]$, so $\tilde{b} \circ \partial^p = Id$. For $p = 1$ we have $\tilde{b}[d\alpha] = [df \wedge (\alpha - \alpha_0)]$ where $\alpha_0 \in \mathbb{C}$ is such that $\alpha|_{X_0} = \alpha_0$. This shows that in degree 1 $\tilde{b}$ gives a canonical splitting of the exact sequence (@).

6.2 $\tilde{\mathcal{A}}$ -structures.

Let us consider now the $\mathbb{C}$ -algebra

$$\tilde{\mathcal{A}} := \left\{ \sum_{\nu \geq 0} P_\nu(a).b^\nu \right\}$$

where $P_\nu \in \mathbb{C}[z]$, and the commutation relation $a.b - b.a = b^2$, assuming that left and right multiplications by a are continuous for the b -adic topology of $\tilde{\mathcal{A}}$. Define the following complexes of sheaves of left $\tilde{\mathcal{A}}$ -modules on X :

$$(\Omega'^\bullet[[b]], D^\bullet) \quad \text{and} \quad (\Omega''^\bullet[[b]], D^\bullet) \quad \text{where} \tag{4}$$

$$\Omega'^\bullet[[b]] := \sum_{j=0}^{+\infty} b^j.\omega_j \quad \text{with} \quad \omega_0 \in \hat{K}^p$$

$$\Omega''^\bullet[[b]] := \sum_{j=0}^{+\infty} b^j.\omega_j \quad \text{with} \quad \omega_0 \in \hat{I}^p$$

$$D\left(\sum_{j=0}^{+\infty} b^j.\omega_j\right) = \sum_{j=0}^{+\infty} b^j.(d\omega_j - df \wedge \omega_{j+1})$$

$$a.\sum_{j=0}^{+\infty} b^j.\omega_j = \sum_{j=0}^{+\infty} b^j.(f.\omega_j + (j-1).\omega_{j-1}) \quad \text{with the convention} \quad \omega_{-1} = 0$$

$$b.\sum_{j=0}^{+\infty} b^j.\omega_j = \sum_{j=1}^{+\infty} b^j.\omega_{j-1}$$

It is easy to check that D is $\tilde{\mathcal{A}}$ -linear and that $D^2 = 0$. We have a natural inclusion of complexes of left $\tilde{\mathcal{A}}$ -modules

$$\tilde{i} : (\Omega''^\bullet[[b]], D^\bullet) \rightarrow (\Omega'^\bullet[[b]], D^\bullet).$$

Remark that we have natural morphisms of complexes

$$u : (\hat{I}^\bullet, d^\bullet) \rightarrow (\Omega''^\bullet[[b]], D^\bullet)$$

$$v : (\hat{K}^\bullet, d^\bullet) \rightarrow (\Omega'^\bullet[[b]], D^\bullet)$$

and that these morphisms are compatible with i . More precisely, this means that we have the commutative diagram of complexes

$$\begin{array}{ccc} (\hat{I}^\bullet, d^\bullet) & \xrightarrow{u} & (\Omega''^\bullet[[b]], D^\bullet) \\ \downarrow i & & \downarrow \tilde{i} \\ (\hat{K}^\bullet, d^\bullet) & \xrightarrow{v} & (\Omega'^\bullet[[b]], D^\bullet) \end{array}$$

The following theorem is a variant of theorem 2.2.1. of [B.II].

Théorème 6.2.1 *Let X be a connected complex manifold of dimension $n + 1$ and $f : X \rightarrow \mathbb{C}$ a non constant holomorphic function such that*

$$\{x \in X / df = 0\} \subset f^{-1}(0).$$

Then the morphisms of complexes u and v introduced above are quasi-isomorphisms. Moreover, the isomorphisms that they induce on the cohomology sheaves of these complexes are compatible with the actions of a and b .

This theorem builds a natural structure of left $\tilde{\mathcal{A}}$ -modules on each of the complex $(\hat{K}^\bullet, d^\bullet)$, $(\hat{I}^\bullet, d^\bullet)$, $([\hat{K}/\hat{I}]^\bullet, d^\bullet)$ and $f^*(\hat{\Omega}_{\mathbb{C}}^1), (\tilde{K}^\bullet, d^\bullet)$ in the derived category of bounded complexes of sheaves of $\mathbb{C}$ -vector spaces on X .

Moreover the short exact sequences

$$\begin{aligned} 0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0 \\ 0 \rightarrow f^*(\hat{\Omega}_{\mathbb{C}}^1) \rightarrow (\hat{K}^\bullet, d^\bullet), (\hat{I}^\bullet, d^\bullet) \rightarrow (\tilde{K}^\bullet, d^\bullet) \rightarrow 0 \end{aligned}$$

are equivalent to short exact sequences of complexes of left $\tilde{\mathcal{A}}$ -modules in the derived category.

PROOF. We have to prove that for any $p \geq 0$ the maps $\mathcal{H}^p(u)$ and $\mathcal{H}^p(v)$ are bijective and compatible with the actions of a and b . The case of $\mathcal{H}^p(v)$ is handled (at least for $n \geq 2$ and f reduced) in prop. 2.3.1. of [B.II]. To seek for completeness and for the convenience of the reader we shall treat here the case of $\mathcal{H}^p(u)$.

First we shall prove the injectivity of $\mathcal{H}^p(u)$. Let $\alpha = df \wedge \beta \in \hat{I}^p \cap \text{Ker } d$ and assume that we can find $U = \sum_{j=0}^{+\infty} b^j \cdot u_j \in \Omega''^{p-1}[[b]]$ with $\alpha = DU$. Then we have the following relations

$$u_0 = df \wedge \zeta, \quad \alpha = du_0 - df \wedge u_1 \quad \text{and} \quad du_j = df \wedge u_{j+1} \quad \forall j \geq 1.$$

For $j \geq 1$ we have $[du_j] = b[du_{j+1}]$ in $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$; using corollary 2.2. of [B.II] which gives the b -separation of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$, this implies $[du_j] = 0, \forall j \geq 1$ in $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$. For instance we can find $\beta_1 \in \hat{K}^{p-1}$ such that $du_1 = d\beta_1$. Now, by de Rham, we can write $u_1 = \beta_1 + d\xi_1$ for $p \geq 2$, where $\xi_1 \in \hat{\Omega}^{p-2}$. Then we conclude that $\alpha = -df \wedge d(\xi_1 + \zeta)$ and $[\alpha] = 0$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$.

For $p = 1$ we have $u_1 = 0$ and $[\alpha] = [-df \wedge d\xi_1] = 0$ in $\mathcal{H}^1(\hat{I}^\bullet, d^\bullet)$.

We shall show now that the image of $\mathcal{H}^p(u)$ is dense in $\mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ for its b -adic topology. Let $\Omega := \sum_{j=0}^{+\infty} b^j \cdot \omega_j \in \Omega''^p[[b]]$ such that $D\Omega = 0$. The following relations holds $d\omega_j = df \wedge \omega_{j+1} \quad \forall j \geq 0$ and $\omega_0 \in \hat{I}^p$. The corollary 2.2. of [B.II] again allows to find $\beta_j \in \hat{K}^{p-1}$ for any $j \geq 0$ such that $d\omega_j = d\beta_j$. Fix $N \in \mathbb{N}^*$. We have

$$D\left(\sum_{j=0}^N b^j \cdot \omega_j\right) = b^N \cdot d\omega_N = D(b^N \cdot \beta_N)$$

and $\Omega_N := \sum_{j=0}^N b^j \omega_j - b^N \beta_N$ is D -closed and in $\Omega''^p[[b]]$. And we have $\Omega - \Omega_N \in b^N \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$, so the sequence $(\Omega_N)_{N \geq 1}$ converges to Ω in $\mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ for its b -adic topology. Let us show that each Ω_N is in the image of $\mathcal{H}^p(u)$.

Write $\Omega_N := \sum_{j=0}^N b^j \omega_j$. The condition $D\Omega_N = 0$ implies $dw_N = 0$ and $dw_{N-1} = df \wedge w_N = 0$. If we write $w_N = dv_N$ we obtain $d(w_{N-1} + df \wedge v_N) = 0$ and $\Omega_N - D(b^N v_N)$ is of degree $N-1$ in b . For $N=1$ we are left with $w_0 + b.w_1 - (-df \wedge v_1 + b.dv_1) = w_0 + df \wedge v_1$ which is in $\hat{I}^p \cap \text{Ker } d$ because $dw_0 = df \wedge dv_1$.

To conclude it is enough to know the following two facts

- i) The fact that $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ is complete for its b -adic topology.
- ii) The fact that $\text{Im}(\mathcal{H}^p(u)) \cap b^N \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet) \subset \text{Im}(\mathcal{H}^p(u) \circ b^N) \quad \forall N \geq 1$.

Let us first conclude the proof of the surjectivity of $\mathcal{H}^p(u)$ assuming i) and ii). For any $[\Omega] \in \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$ we know that there exists a sequence $(\alpha_N)_{N \geq 1}$ in $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ with $\Omega - \mathcal{H}^p(u)(\alpha_N) \in b^N \mathcal{H}^p(\Omega''^\bullet[[b]], D^\bullet)$. Now the property ii) implies that we may choose the sequence $(\alpha_N)_{N \geq 1}$ such that $[\alpha_{N+1}] - [\alpha_N]$ lies in $b^N \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. So the property i) implies that the Cauchy sequence $([\alpha_N])_{N \geq 1}$ converges to $[\alpha] \in \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. Then the continuity of $\mathcal{H}^p(u)$ for the b -adic topologies coming from its b -linearity, implies $\mathcal{H}^p(u)([\alpha]) = [\Omega]$.

The compatibility with a and b of the maps $\mathcal{H}^p(u)$ and $\mathcal{H}^p(v)$ is an easy exercise.

Let us now prove properties i) and ii).

The property i) is a direct consequence of the completion of $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ for its b -adic topology given by the corollary 2.2. of [B.II] and the b -linear isomorphism $\tilde{b}$ between $\mathcal{H}^p(\hat{K}^\bullet, d^\bullet)$ and $\mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$ constructed in the lemma 2.1.1. above.

To prove ii) let $\alpha \in \hat{I}^p \cap \text{Ker } d$ and $N \geq 1$ such that

$$\alpha = b^N \Omega + DU$$

where $\Omega \in \Omega''^p[[b]]$ satisfies $D\Omega = 0$ and where $U \in \Omega''^{p-1}[[b]]$. With obvious notations we have

$$\begin{aligned} \alpha &= du_0 - df \wedge u_1 \\ &\dots \\ 0 &= du_j - df \wedge u_{j+1} \quad \forall j \in [1, N-1] \\ &\dots \\ 0 &= \omega_0 + du_N - df \wedge u_{N+1} \end{aligned}$$

which implies $D(u_0 + b.u_1 + \dots + b^N.u_N) = \alpha + b^N.du_N$ and the fact that du_N lies in $\hat{I}^p \cap \text{Ker } d$. So we conclude that $[\alpha] + b^N.[du_N]$ is in the kernel of $\mathcal{H}^p(u)$ which is 0. Then $[\alpha] \in b^N \mathcal{H}^p(\hat{I}^\bullet, d^\bullet)$. ■

REMARK. The map

$$\beta : (\Omega'[[b]]^\bullet, D^\bullet) \rightarrow (\Omega''[[b]]^\bullet, D^\bullet)$$

defined by $\beta(\Omega) = b.\Omega$ commutes to the differentials and with the action of b . It induces the isomorphism $\tilde{b}$ of the lemma 6.1.1 on the cohomology sheaves. So it is a quasi-isomorphism of complexes of $\mathbb{C}[[b]]$ -modules.

To prove this fact, it is enough to verify that the diagram

$$\begin{array}{ccc} (\hat{K}^\bullet, d^\bullet) & \xrightarrow{v} & (\Omega'[[b]]^\bullet, D^\bullet) \\ \downarrow \tilde{b} & & \downarrow \beta \\ (\hat{I}^\bullet, d^\bullet) & \xrightarrow{u} & (\Omega''[[b]]^\bullet, D^\bullet) \end{array}$$

induces commutative diagrams on the cohomology sheaves.

But this is clear because if $\alpha = d\xi$ lies in $\hat{K}^p \cap \text{Ker } d$ we have $D(b.\xi) = b.d\xi - df \wedge \xi$ so $\mathcal{H}^p(\beta) \circ \mathcal{H}^p(v)([\alpha]) = \mathcal{H}^p(u) \circ \mathcal{H}^p(\tilde{b})([\alpha])$ in $\mathcal{H}^p(\Omega''[[b]]^\bullet, D^\bullet)$. ■

6.3 The finiteness theorem.

Let us recall some basic definitions on the left modules over the algebra $\tilde{\mathcal{A}}$.

Now let E be any left $\tilde{\mathcal{A}}$ -module, and define $B(E)$ as the b -torsion of E . that is to say

$$B(E) := \{x \in E \mid \exists N \quad b^N.x = 0\}.$$

Define $A(E)$ as the a -torsion of E and

$$\hat{A}(E) := \{x \in E \mid \mathbb{C}[[b]].x \subset A(E)\}.$$

Remark that $B(E)$ and $\hat{A}(E)$ are sub- $\tilde{\mathcal{A}}$ -modules of E but that $A(E)$ is not stable by b .

Définition 6.3.1 A left $\tilde{\mathcal{A}}$ -module E is called **small** when the following conditions hold

1. E is a finite type $\mathbb{C}[[b]]$ -module ;
2. $B(E) \subset \hat{A}(E)$;
3. $\exists N \mid a^N.\hat{A}(E) = 0$;

Recall that for E small we have always the equality $B(E) = \hat{A}(E)$ (see [B.I] lemme 2.1.2) and that this complex vector space is finite dimensional. The quotient $E/B(E)$ is an (a,b) -module called **the associate (a,b) -module** to E .

Conversely, any left $\tilde{\mathcal{A}}$ -module E such that $B(E)$ is a finite dimensional $\mathbb{C}$ -vector space and such that $E/B(E)$ is an (a,b) -module is small.

The following easy criterium to be small will be used later :

Lemme 6.3.2 *A left $\tilde{\mathcal{A}}$ -module E is small if and only if the following conditions hold :*

1. $\exists N \ / \ a^N \cdot \hat{A}(E) = 0$;
2. $B(E) \subset \hat{A}(E)$;
3. $\cap_{m \geq 0} b^m \cdot E \subset \hat{A}(E)$;
4. $\text{Ker } b$ and $\text{Coker } b$ are finite dimensional complex vector spaces.

As the condition 3 in the previous lemma has been omitted in [B.II] (but this does not affect this article because this lemma was used only in a case where this condition 3 was satisfied, thanks to proposition 2.2.1. of *loc. cit.*), we shall give the (easy) proof.

PROOF. First the conditions 1 to 4 are obviously necessary. Conversely, assume that E satisfies these four conditions. Then condition 2 implies that the action of b on $\hat{A}(E)/B(E)$ is injective. But the condition 1 implies that $b^{2N} = 0$ on $\hat{A}(E)$ (see [B.I]). So we conclude that $\hat{A}(E) = B(E) \subset \text{Ker } b^{2N}$ which is a finite dimensional complex vector space using condition 4 and an easy induction. Now $E/B(E)$ is a $\mathbb{C}[[b]]$ -module which is separated for its b -adic topology. The finiteness of $\text{Coker } b$ now shows that it is a free finite type $\mathbb{C}[[b]]$ -module concluding the proof. ■

Définition 6.3.3 *We shall say that a left $\tilde{\mathcal{A}}$ -module E is **geometric** when E is small and when it associated (a,b) -module $E/B(E)$ is geometric.*

The main result of this section is the following theorem, which shows that the Gauss-Manin connection of a proper holomorphic function produces geometric $\tilde{\mathcal{A}}$ -modules associated to vanishing cycles and nearby cycles.

Théorème 6.3.4 *Let X be a connected complex manifold of dimension $n + 1$ where n is a natural integer, and let $f : X \rightarrow D$ be an non constant proper holomorphic function on an open disc D in $\mathbb{C}$ with center 0. Let us assume that df is nowhere vanishing outside of $X_0 := f^{-1}(0)$. Then the $\tilde{\mathcal{A}}$ -modules*

$$\mathbb{H}^j(X, (\hat{K}^\bullet, d^\bullet)) \quad \text{and} \quad \mathbb{H}^j(X, (\hat{I}^\bullet, d^\bullet))$$

are geometric for any $j \geq 0$.

In the proof we shall use the $\mathcal{C}^\infty$ version of the complex $(\hat{K}^\bullet, d^\bullet)$. We define K_∞^p as the kernel of $\wedge df : \mathcal{C}^{\infty,p} \rightarrow \mathcal{C}^{\infty,p+1}$ where $\mathcal{C}^{\infty,j}$ denote the sheaf of $\mathcal{C}^\infty$ -forms on X of degree p , let $\hat{K}_\infty^p$ be the f -completion and $(\hat{K}_\infty^\bullet, d^\bullet)$ the corresponding de Rham complex.

The next lemma is proved in [B.II] (lemma 6.1.1.)

Lemme 6.3.5 *The natural inclusion*

$$(\hat{K}^\bullet, d^\bullet) \hookrightarrow (\hat{K}_\infty^\bullet, d^\bullet)$$

induce a quasi-isomorphism.

REMARK. As the sheaves $\hat{K}_\infty^\bullet$ are fine, we have a natural isomorphism

$$\mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \simeq H^p(\Gamma(X, \hat{K}_\infty^\bullet), d^\bullet).$$

Let us denote by X_1 the generic fiber of f . Then X_1 is a smooth compact complex manifold of dimension n and the restriction of f to $f^{-1}(D^*)$ is a locally trivial $\mathcal{C}^\infty$ bundle with typical fiber X_1 on $D^* = D \setminus \{0\}$, if the disc D is small enough around 0. Fix now $\gamma \in H_p(X_1, \mathbb{C})$ and let $(\gamma_s)_{s \in D^*}$ the corresponding multivalued horizontal family of p -cycles $\gamma_s \in H_p(X_s, \mathbb{C})$. Then, for $\omega \in \Gamma(X, \hat{K}_\infty^p \cap \text{Ker } d)$, define the multivalued holomorphic function

$$F_\omega(s) := \int_{\gamma_s} \frac{\omega}{df}.$$

Let now

$$\Xi := \bigoplus_{\alpha \in \mathbb{Q} \cap]-1, 0], j \in [0, n]} \mathbb{C}[[s]] \cdot s^\alpha \cdot \frac{(\text{Log } s)^j}{j!}.$$

This is an $\tilde{\mathcal{A}}$ -modules with a acting as multiplication by s and b as the primitive in s without constant. Now if $\hat{F}_\omega$ is the asymptotic expansion at 0 of F_ω , it is an element in Ξ , and we obtain in this way an $\tilde{\mathcal{A}}$ -linear map

$$\text{Int} : \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \rightarrow H^p(X_1, \mathbb{C}) \otimes_{\mathbb{C}} \Xi.$$

To simplify notations, let $E := \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet))$. Now using Grothendieck theorem [G.65], there exists $N \in \mathbb{N}$ such that $\text{Int}(\omega) \equiv 0$, implies $a^N \cdot [\omega] = 0$ in E . As the converse is clear we conclude that $\hat{A}(E) = \text{Ker}(\text{Int})$. It is also clear that $B(E) \subset \text{Ker}(\text{Int})$ because Ξ has no b -torsion. So we conclude that E satisfies properties 1 and 2 of the lemma 6.3.2. The property 3 is also true because of the regularity of the Gauss-Manin connection of f .

END OF THE PROOF OF THEOREM 6.3.4. To show that $E := \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet))$ is small, it is enough to prove that E satisfies the condition 4 of the lemma 6.3.2. Consider now the long exact sequence of hypercohomology of the exact sequence of complexes

$$0 \rightarrow (\hat{I}^\bullet, d^\bullet) \rightarrow (\hat{K}^\bullet, d^\bullet) \rightarrow ([\hat{K}/\hat{I}]^\bullet, d^\bullet) \rightarrow 0.$$

It contains the exact sequence

$$\mathbb{H}^{p-1}(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet)) \rightarrow \mathbb{H}^p(X, (\hat{I}^\bullet, d^\bullet)) \xrightarrow{\mathbb{H}^p(i)} \mathbb{H}^p(X, (\hat{K}^\bullet, d^\bullet)) \rightarrow \mathbb{H}^p(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$$

and we know that b is induced on the complex of $\tilde{\mathcal{A}}$ -modules quasi-isomorphic to $(\hat{K}^\bullet, d^\bullet)$ by the composition $i \circ \tilde{b}$ where $\tilde{b}$ is a quasi-isomorphism of complexes of $\mathbb{C}[[b]]$ -modules. This implies that the kernel and the cokernel of $\mathbb{H}^p(i)$ are isomorphic (as $\mathbb{C}$ -vector spaces) to $\text{Ker } b$ and $\text{Coker } b$ respectively. Now to prove that E satisfies condition 4 of the lemma 6.3.2 it is enough to prove finite dimensionality for the vector spaces $\mathbb{H}^j(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$ for all $j \geq 0$.

But the sheaves $[\hat{K}/\hat{I}]^j \simeq [\text{Ker } df / \text{Im } df]^j$ are coherent on X and supported in X_0 . The spectral sequence

$$E_2^{p,q} := H^q(H^p(X, [\hat{K}/\hat{I}]^\bullet, d^\bullet))$$

which converges to $\mathbb{H}^j(X, ([\hat{K}/\hat{I}]^\bullet, d^\bullet))$, is a bounded complex of finite dimensional vector spaces by Cartan-Serre. This gives the desired finite dimensionality.

To conclude the proof, we want to show that $E/B(E)$ is geometric. But this is an easy consequence of the regularity of the Gauss-Manin connexion of f and of the Monodromy theorem, which are already incoded in the definition of Ξ : the injectivity on $E/B(E)$ of the $\tilde{\mathcal{A}}$ -linear map Int implies that $E/B(E)$ is geometric. Remark now that the piece of exact sequence above gives also the fact that $\mathbb{H}^p(X, (\hat{I}^\bullet, d^\bullet))$ is geometric, because it is an exact sequence of $\tilde{\mathcal{A}}$ -modules. ■

REMARK. It is easy to see that the properness assumption on f is only used for two purposes :

- To have a (global) $\mathcal{C}^\infty$ Milnor fibration on a small punctured disc around 0, with a finite dimensional cohomology for the Milnor fiber.
- To have compactness of the singular set $\{df = 0\}$.

This allows to give with the same proof an analogous finiteness result in many other situations.

Bibliography

- [Br.70] Brieskorn, E. *Die Monodromie der Isolierten Singularitäten von Hyperflächen*, Manuscripta Math. 2 (1970), p. 103-161.
- [B.84] Barlet, D. *Contribution du cup-produit de la fibre de Milnor aux pôles de $|f|^{2\lambda}$* , Ann. Inst. Fourier (Grenoble) t. 34, fasc. 4 (1984), p. 75-107.
- [B. 93] Barlet, D. *Théorie des (a,b) -modules I*, in Complex Analysis and Geometry, Plenum Press, (1993), p. 1-43.
- [B. 95] Barlet, D. *Théorie des (a,b) -modules II. Extensions*, in Complex Analysis and Geometry, Pitman Research Notes in Mathematics Series 366 Longman

(1997), p. 19-59.

- [B. 05] Barlet, D. *Module de Brieskorn et forme hermitiennes pour une singularité isolée d'hypersurface*, revue de l'Inst. E. Cartan (Nancy) 18 (2005), p. 19-46.
- [B.I] Barlet, D. *Sur certaines singularités non isolées d'hypersurfaces I*, Bull. Soc. math. France 134 (2), 2006, p.173-200.
- [B. II] Barlet, D. *Sur certaines singularités d'hypersurfaces II*, J. Alg. Geom. 17 (2008), p. 199-254.
- [B. 08] Barlet, D. *Two finiteness theorem for regular (a,b) -modules*, preprint Institut E. Cartan (Nancy) (2008) n^05 , p. 1-38, arXiv:0801.4320 (math. AG and math. CV)
- [B.09] Barlet,D. *Périodes évanescentes et (a,b) -modules monogènes*, Bollettino U.M.I. (9) II (2009) p.651-697.
- [B.III] Barlet, D. *Sur les fonctions à lieu singulier de dimension 1* , Bull. Soc. math. France 137 (4), 2009, p.587-612.
- [B.10] Barlet,D. *Le thème d'une période évanescence*, preprint Institut E. Cartan (Nancy) (2009) n^033 , p.1-57.
- [B.-S. 04] Barlet, D. et Saito, M. *Brieskorn modules and Gauss-Manin systems for non isolated hypersurface singularities*, J. Lond. Math. Soc. (2) 76 (2007) n^01 p. 211-224.
- [Bj.93] Björk, J-E, *Analytic D-modules and applications*, Kluwer Academic publishers (1993).
- [K.76] Kashiwara, M. *b-function and holonomic systems*, Inv. Math. 38 (1976) p. 33-53.
- [M.74] Malgrange, B. *Intégrale asymptotique et monodromie*, Ann. Sc. Ec. Norm. Sup. 7 (1974), p.405-430.

- [M. 75] Malgrange, B. *Le polynôme de Bernstein d'une singularité isolée*, in Lect. Notes in Math. 459, Springer (1975), p.98-119.
- [S. 89] Saito, M. *On the structure of Brieskorn lattices*, Ann. Inst. Fourier 39 (1989), p.27-72.